

BEHAVIOUR

The subtle difference between the sexes

DOI:
10.1038/nrn2226

How is sexual dimorphism in behaviour represented in the brain? Numerous attempts to identify a unique brain structure or circuit that underlies sex-specific traits within one species have so far failed. Dulac and colleagues now show that vomeronasal neurons that express the *Trpc2* ion channel are important for female-specific traits in mating and social behaviour in mice.

The current hypothesis regarding the development of sex-specific traits claims that female-specific brain centres form by default. In males, male hormones stimulate the formation of male-specific brain centres and prevent the formation of female-specific centres. The authors

investigated sexual and social behaviour in female mice that lacked *Trpc2* (*Trpc2*^{-/-} mice) — an ion channel that is implicated in pheromone perception and is exclusively expressed in the vomeronasal organ (VNO) — or in which the VNO had been surgically removed (VNOx mice).

The lack of expression of *Trpc2* caused females to display male sexual and courtship behaviour towards both sexes. The behaviour of these female mice was indistinguishable from that of male mice that lacked one or both *Trpc2* alleles. These mice also lost the ability to distinguish between different genders, but continued to display male-like sexual behaviour traits. Furthermore, the behaviour of *Trpc2*^{-/-} females was similar throughout their oestrous cycle. Therefore, the authors concluded that the behaviour of *Trpc2*^{-/-} females is the result of the expression of a male-like trait.

To determine whether sexual traits are established during development or are subject to adult plasticity, the authors surgically removed the VNO from adult animals. VNOx males displayed male-like sexual and courtship behaviour towards both females and males. Thus, they behaved like *Trpc2*^{-/-} and *Trpc2*^{+/-} males, which also no longer discriminated between the sexes. Removal of the VNO in adult females reversed their sexual behaviour, as they now

showed male-like sexual display that was similar to that shown by *Trpc2*^{-/-} females. The authors concluded that sex-specific behavioural traits do not stem from the formation of specific circuits during development, but that neural circuits that are targeted by VNO neurons are determinants for sex-specific behaviour.

In accordance with the observed male-like traits of *Trpc2*^{-/-} and VNOx female mice, the authors also reported that maternal behaviour was largely repressed in these animals. Importantly, no significant changes in hormone levels were detected when *Trpc2*^{-/-} and VNOx female mice were compared with wild-type females.

In summary, female mice that do not express *Trpc2* or that lack the VNO display male-like sexual and courtship behaviour and have reduced maternal instincts. These findings indicate that the VNO has a role in suppressing male-like behaviour in females. Based on their findings, the authors proposed a new hypothesis: that during development, both sexes form male- and female-trait centres in their brains. In adult females, pheromones activate the female-trait centres and inhibit the activity of the male centres, resulting in female-specific sexual behaviour. Similarly, pheromones in males specifically inhibit the activity of female centres and activate male centres. It remains to be determined whether *Trpc2*^{-/-} or VNOx males display female traits.

Claudia Wiedemann

ORIGINAL RESEARCH PAPER Kimchi, T., Xu, J. & Dulac, C. A functional circuit underlying male sexual behaviour in the female mouse brain. *Nature* 05 Aug 2007 (doi:10.1038/nature06089)