


# Touching




**Ehud Ahissar**

# Touching

- Body-world interface
- Mechanisms of sensory processing (across senses)
- Motor-sensory coupling
- Passive vs active touch
- Neuronal coding
- Morphological coding


# Body-world interface

## Underneath the skin




**Figure 8.3** The skin harbors a variety of morphologically distinct mechanoreceptors. This diagram represents the smooth, hairless (also called glabrous) skin of the fingertip. The major characteristics of the various receptor types are summarized in Table 8.1. (After Darian-Smith, 1984.)



# Mechanoreception underneath the skin



~200  $\mu\text{m}$

# Mechanoreception underneath the skin





# Proprioceptive receptor types

Name:

Muscle spindle receptors

Golgi tendon organs


Joint receptors

Sensitive to:

muscle length change

muscle tension

Joint angle



# Body-world interface

## Underneath the skin

**TABLE 8.1**  
The Major Classes of Somatic Sensory Receptors

Receptor type	Anatomical characteristics	Associated axons <sup>a</sup> (and diameters)	Axonal conduction velocities	Location	Function	Rate of adaptation	Threshold of activation
Free nerve endings	Minimally specialized nerve endings	A $\delta$ C	2–20 m/s .5 – 2 m/s	All skin	Pain, temperature, crude touch	Slow	High
Meissner's corpuscles	Encapsulated; between dermal papillae	A $\beta$ 6–12 $\mu$ m	30 – 70 m/s	Principally glabrous skin	Touch, pressure (dynamic)	Rapid	Low
Pacinian corpuscles	Encapsulated; onionlike covering	A $\beta$ 6–12 $\mu$ m		Subcutaneous tissue, interosseous membranes, viscera	Deep pressure, vibration (dynamic)	Rapid	Low
Merkel's disks	Encapsulated; associated with peptide-releasing cells	A $\beta$		All skin, hair follicles	Touch, pressure (static)	Slow	Low
Ruffini's corpuscles	Encapsulated; oriented along stretch lines	A $\beta$ 6–12 $\mu$ m		All skin	Stretching of skin	Slow	Low
Muscle spindles	Highly specialized (see Figure 8.5 and Chapter 15)	Ia and II	80 – 120 m/s	Muscles	Muscle length	Both slow and rapid	Low
Golgi tendon organs	Highly specialized (see Chapter 15)	Ib	80 – 120 m/s	Tendons	Muscle tension	Slow	Low
Joint receptors	Minimally specialized	—		Joints	Joint position	Rapid	Low

Mechano-receptors  
(ex-afferents)


Proprio-(re)ceptors  
(re-afferents)

<sup>a</sup>In the 1920s and 1930s, there was a virtual cottage industry classifying axons according to their conduction velocity. Three main categories were discerned, called




# Receptors

## Evolutionary specialization




(a) Free nerve endings



(b) Free nerve endings of root hair plexus




(c) Merkel cells and Merkel's discs




Selection process - ?


Morphological processing




(f) Ruffini corpuscle



Layers of collagen fibers separated by fluid



(e) Pacinian corpuscle



(d) Meissner's corpuscle


Dermis

# Signal transduction


# Transduction

The receptor potential is produced by a **mechanically sensitive channel** that opens when the membrane is deformed

The channel is permeable to positive ions, primarily  $\text{Na}^+$ ,  $\text{K}^+$  and  $\text{Ca}^{2+}$



# Transduction




**Figure 21-2** Mechanoreceptors are depolarized by stretch of the cell membrane and the depolarization is proportional to the stimulus amplitude.

**A.** The spindle organ in skeletal muscle mediates limb proprioception. These receptors signal muscle length and the speed at which the

# Transduction


Mechanically activated currents in **Merkel cells** depend on **Piezo2**.



**Receptive Fields (RFs):**


**Spatial and temporal**

# Receptive Fields (RFs): Spatial and temporal



RF size?




# Receptive Fields (RFs): Spatial and temporal




Response dynamics?




# Receptive Fields (RFs): Spatial and temporal



# Cutaneous Mechanoreceptor Channels




# Receptor density



**Figure 22-4 The distribution of receptor types in the human hand varies.** The number of sensory nerve fibers innervating an area is indicated by the stippling density, with the highest density of receptors shown by the heaviest stippling. (**RA** = 5 rapidly adapting, **SA** = 5 slowly adapting.) Meissner's corpuscles (**RA**) and Merkel disk receptors (**SA I**) are the most numerous receptors; they are distributed preferentially on the distal half of the fingertip. Pacinian corpuscles (**PC**) and Ruffini endings (**SA II**) are much less common; they are distributed more uniformly on the hand, showing little differentiation of the distal and proximal regions. The fingertips are the most densely innervated region of skin in the human body, receiving approximately 300 mechanoreceptive nerve fibers per square centimeter. The number of mechanoreceptive fibers is reduced to 120/cm<sup>2</sup> in the proximal phalanges, and to 50/cm<sup>2</sup> in the palm. (Adapted from Vallbo and Johansson 1978.)

# Neurometric - psychometric matching


## Spatial resolution (by JND)




- Break ?

# Signal conduction

# Sensory signal conduction



# Sensory signal conduction






# Sensory signal conduction

TABLE 8.1


The Major Classes of Somatic Sensory Receptors


Receptor type	Anatomical characteristics	Associated axons <sup>a</sup> (and diameters)
Free nerve endings	Minimally specialized nerve endings	C, A $\delta$ , A $\beta$ , C
Meissner's corpuscles	Encapsulated; between dermal papillae	A $\beta$ 6–12 $\mu$ m
Pacinian corpuscles	Encapsulated; onionlike covering	A $\beta$ 6–12 $\mu$ m
Merkel's disks	Encapsulated; associated with peptide-releasing cells	A $\beta$
Ruffini's corpuscles	Encapsulated; oriented along stretch lines	A $\beta$ 6–12 $\mu$ m
Muscle spindles	Highly specialized (see Figure 8.5 and Chapter 15)	Ia and II
Golgi tendon organs	Highly specialized (see Chapter 15)	Ib
Joint receptors	Minimally specialized	—



<sup>a</sup>In the 1920s and 1930s, there was a virtual cottage industry classifying axons according to their conduction velocity. Three main categories were discerned, called

# Sensory signal conduction








**Figure 8.7** Diagram of the somatic sensory portions of the thalamus and their cortical targets in the postcentral gyrus. The ventral posterior nuclear complex comprises the VPM, which relays somatic sensory information carried by the trigeminal system from the face, and the VPL, which relays somatic sensory information from the rest of the body. Inset above shows organization of the primary somatosensory cortex in the postcentral gyrus, shown here in a section cutting across the gyrus from anterior to posterior. (After Brodal, 1992, and Jones et al., 1982.)

# The Homunculi

A Sensory homunculus




B Motor homunculus



# The Homunculi

Relative size reflects innervation density

phylogenetically



# The Homunculi

## Accurate spatial organization





Figure 23-9 The representation of whiskers in the somatosensory cortex of the rat. (Adapted from [Bennett-Clarke et al. 1997](#)).


# The Homunculi

Relative size reflects innervation density

ontogenetically



# Motor signal conduction



## The cortico-spinal tract

30% - M1

30% - premotor



30% - somatosensory, parietal

## The cortico-bulbar tract

Face, head, neck



# Motor signal conduction



The cortico-spinal tract

(not reversal of the afferent pathway)


# Motor signal conduction

Midbrain




Cerebral peduncle

Pons




The cortico-spinal tract

Medulla



Cortico-centric view  
Vs  
Evolution-based view

Medulla-spinal cord junction




Lateral corticospinal tract

Lateral column

Cervical spinal cord

1<sup>st</sup> synapse




Lateral intermediate zone and lateral motor nuclei



# Sensory signal conduction



## The vibrissal system





# Sensory signal conduction

## The vibrissal system


whisker



**b**



b



More on this in the  
Active-Sensing lecture

- Lemniscal
- Extralemniscal
- Paralemniscal

Follicle/whisker  
complex

# Common mechanisms of sensory processing



# Rich muscular system








Fig. 5. Medial aspect of two neighbouring follicles C<sub>1</sub> and C<sub>2</sub> and their motor and sensory nerves. Right whisker tip. The medial sacculus of the follicular muscle (M) belonging to follicle C<sub>2</sub> was divided and reflected to allow the inter-follicular space. Two longitudinal branches of the facial nerve (V7) send bifurcating nerves to the follicular muscle. The row nerve (F3) gives off the follicular nerves (F) perforating the postero-medial aspect of the follicles, and skin here give off the nerves (F-2) reaching the dermis and sending the coin nerve (S.A.) to the central body of the follicle.

# Receptor types

eye




R G B

finger

RAI

SAI

RAII



RA

SA

PC

Ruffini endings

whisker





Figure 1


eye

@ 1°



# Receptors mix in clusters



whisker



finger



Merkel cells

## Receptor convergence / divergence

**Human eye:** 5M cones (+ 120M rods) --> 1M fibers

**Human skin:** 2,500 receptors/cm<sup>2</sup> --> 300 fibers / cm<sup>2</sup>

**Rat whisker:** 2,000 receptors --> 300 fibers


~ 10 -> 1 convergence

**Human ear:** 3,000 hair cells --> 30,000 fibers

~ 1 -> 10 divergence

# Processing stations


eye



Receptors

Bipolar cells


Ganglion cells



Thalamus

Cortex





finger



Receptors

Ganglion cells


Brainstem cells



Thalamus

Cortex





whisker



Receptors

Ganglion cells

Brainstem cells



Thalamus

Cortex






# Spatial processing (by Lateral inhibition)

eye


finger

?

whisker



Drive for clustering?



# Efficient coding

## (by coding changes only)

### Changes in time:

- Intrinsic in individual neurons
- Starting at the receptor level


### Changes in space:

- Circuits of neurons
- Starting after lateral inhibition


# Temporal filtering (by intrinsic factors)

eye

whisker




finger





# Neurometric - psychometric matching

sensitivity



**Figure 22-6B** The threshold for detecting vibration corresponds to the tuning threshold of the mechanoreceptor. The sensitivity threshold for Meissner's corpuscles is lowest for frequencies of 20-50 Hz. Pacinian corpuscles sense higher frequencies. (Adapted from Mountcastle et al. 1972.)

- Break ?

# Passive and active touch

## Passive touch

- Perceptual processing follows sensory events

## Active touch

- Perceptual processing surrounds sensory events:
  - The brain probes the world
  - Compares sensory data with internal expectations
  - Updates internal expectations



## Active touch is done in a loop:

- Change of expectations => probing the world
- probing the world => Change of expectations

## Passive and active touch

### **Passive touch**

- low thresholds
- poor accuracy


### **Active touch**

- higher thresholds
- high accuracy

# Passive and active touch

## Passive touch


- low thresholds
- poor accuracy



Detection

## Active touch

- higher thresholds
- high accuracy



Exploration

Object localization

Object identification

# Passive and active touch

## Passive touch

- low thresholds
- poor accuracy

## Active touch

- higher thresholds
- high accuracy

### Potential underlying mechanism: “Gating”

- Arousal, preparatory, or motor commands “gate out” sensory signals
- Example: Thalamic gating (Sherman & Guillery, JNP. 1996)

Thalamic neurons have 2 modes:

- in drowsiness: hyperpolarized, bursting, low threshold
- in alertness: depolarized, single spikes, high threshold

# Passive and active touch

## **Passive touch**

- low thresholds
- poor accuracy

## **Active touch**

- higher thresholds
- high accuracy

### Underlying mechanisms:

- Additional information
  - expectations
  - accumulation of sensory data over time
  - more coding dimensions
  - increased resolution due to scanning
- close-loop operation

# Passive and active touch

## **Passive touch**

- low thresholds
- poor accuracy

## **Active touch**

- higher thresholds
- high accuracy

### Underlying active mechanisms:

- Additional information
  - expectations
  - accumulation of sensory data over time
  - more coding dimensions
  - increased resolution due to scanning
- close-loop operation




Sensory encoding:

What receptors tell the brain


Sensory organs consist of **receptor arrays**:

**somatosensation**



~200 μm

*Finger pad*




**audition**




10 μm

*cochlea*




**vision**



10 μm

*retina*




Sensory encoding:

What receptors tell the brain

Sensory organs consist of **receptor arrays**:

**somatosensation**



~200  $\mu\text{m}$

*Finger pad*


**audition**



10  $\mu\text{m}$

*cochlea*

**vision**



10  $\mu\text{m}$

*retina*


**Spatial organization => Spatial coding** (“*which* receptors are activated”)



**How neurons encode external events in space?**


## The “labeled-line code”.

a binary code, reporting yes/no about the occurrence of a given event.



Every neuron has a “label”

## Reading out the labeled line code



*reading algorithm:* a location X is pressed if and only if neuron x fires

On what condition will this algorithm be valid?

(X)  ~~$X \Rightarrow$~~   Neuron x fires ~~if and only if~~ X is pressed

**Is this assumption valid?**

**1. The problem of background activity**

## 2. The “problem” of sensor movements

receptors are sensitive to changes

Thus

If both objects and sensors are passive (stationary),  
nothing will be sensed


Thus

Sensors must move in order to sense stationary  
objects

**How sensor motion constrains sensory coding?**



## Reading out the labeled line code



*reading algorithm:* a location X is pressed if and only if neuron x fires

On what condition will this algorithm be valid?

(X)  ~~$\bullet \Rightarrow$~~   $\bullet_x$  —|—|—| Neuron x fires ~~if and only if~~ X is pressed

**Is this assumption valid?**


## 2. The “problem” of sensor motion

sensory encoding:

What receptors tell the brain

Sensory organs consist of **receptor arrays**:

**somatosensation**




~50  $\mu\text{m}$

*Finger pad*

**audition**



**vision**



*retina*


More on this in the  
**Active-Sensing lecture**

**Spatial organization** => **Spatial coding** (“*which* receptors are activated”)


**Movements** => **Temporal coding** (“*when* receptors activated”)

# Receptors

## Evolutionary specialization




(a) Free nerve endings



(b) Free nerve endings of root hair plexus




(c) Merkel cells and Merkel's discs





Selection process - ?


Morphological processing



(f) Ruffini corpuscle




(d) Meissner's corpuscle





(e) Pacinian corpuscle

# Morphological processing




## Morphological phase plane





# Morphological processing

Motor-sensory phase plane  
Morphological phase plane




Sensory



Motor


# Organism-environment attractors



# Touching

- Body-world interface
- Mechanisms of sensory processing (across senses)
- Motor-sensory coupling
- Passive vs active touch
- Neuronal coding
- Morphological coding

# Touching



**The End**