

# Quantum Coherence Effects in Biological Systems — A New “Twist”

R. J. Dwayne Miller

Max Planck Research Group for Structural  
Dynamics

Department of Physics, University of Hamburg,  
CFEL/DESY and  
Departments of Chemistry and Physics,  
University of Toronto



# Acknowledgements

Valentyn Prokorenko MPG,  
**University of Hamburg**

Alexei Halpin (Physics)  
Philip Johnson (Chemistry)  
Oliver Ernst/group  
(Biochem)  
**University of Toronto**

Stephen A. Waschuk  
Leonid S. Brown  
Dept. of Physics  
**University of Guelph**




Ray Gao  
Hubert Jean-Ruel  
Cheng Lu  
Nelson Lui  
German Sciani  
Gustavo Morienna  
**University of Toronto**

Shin-ya Koshihara  
Ken Onda  
**Tokyo Institute of Tech**  
Hideki Yamochi  
**Kyoto University**



# In the beginning....



**Fundamental issue:** how well is the biosystem optimized to this “light food” ??

# Quantum Coherence and Biology

Tenet: Biological systems (at the molecular level)  
have evolved to control the transition state region  
Barrier Crossings (transition state processes)  
occur over atomic length scales  $\Rightarrow$  wave properties  
of matter become significant


*Has Nature evolved to even exploit phase?*

**Coherence properties of waves require an  
interferometer to measure  $\Rightarrow$  Coherent Control  $\equiv$   
Molecular Frame of Reference Interferometer**


Intrepid Surfer Analogy

- Structures of PS II and PS I (protein not shown)

X-Ray structure of Photosystem II from *S. elongatus* with 3.8 Å resolution, file # 1FE1 in PDB.  
A. Zouni et al. *Nature*, 409, 739 (2001).






X-Ray structure of Photosystem I from *S. elongatus* with 2.5 Å resolution, file # 1JBO in PDB.  
P. Jordan et al. *Nature*, 411, 909 (2001).



**Nature is optimized at the coherence limits of mol. responses**

# THE DREAM – CONTROLLING ISOMERIZATION IN RHODOPSIN





<http://users.rcn.com/jkimball.ma.ultranet/BiologyPages/V/Vision.html>

**Ideal System:** Biologically Relevant Photoinduced Function THAT is fast enough to compete with Quantum Decoherence  $\Rightarrow$  **Must Demonstrate under Weak Field Control**


Q. Wang, R. Schonlein, L. Peteanu, R. Mathies, C. Shank, Science 1994, 266, 422

# Bacteriorhodopsin-a Precursor to Rhodopsin



<http://www.science.siu.edu/microbiolog/micr425/Halobacteria96>

## •Bacteriorhodopsin – the smallest chameleon in Nature





<http://www.science.siu.edu/microbiolog/micr425/Halobacteria96>




*all-trans* form: light-adapted ground state →

**Efficiency of isomerization ~ 65%**



⇒ **Reaction Dynamics can be described within a power spectrum**  
re: dominant mode couplings




# Relevant Experimental Work



Wohlleben et al. ChemPhysChem, 6, 850, 2005 (Herek and Motkus groups)



Nuemberger et al. JCP, 125, 044512 (2006)/Vogt et al PRL 2005




Vogt et al. CPL 433, 211 (2006)

⇒ Degree of Control Increases under Strong Field/High Intensity Conditions

# Coherent Control in the Weak Field Limit


## CLOSED QUANTUM SYSTEMS

-no coupling to bath



Single state case:  
1 eigenstate  $\Rightarrow$  1 pathway  
• no interference

$\rightarrow$  no control




bichromic case: 2 pathways  
• linear  
• fixed phase leads to interference

$\rightarrow$  coherent control

## OPEN QUANTUM SYSTEMS


-coupling to bath, or surroundings



- several pathways
- interference at CI
- phase sensitive relaxation/dissipation to bath

$\rightarrow$  coherent control


# Coherent control setup




# Coherent Control of Retinal Photoisomerization\* — Quantum Control of a Biological Function

## GOAL

- Control isomerization efficiency under these restrictions:
  - a) weak field excitation (within **linear response** regime)
  - b) **fixed number of absorbed photons** per laser shot



from: Kobyashi et al., Nature, v. 414 (2001)




from: Tretyak et al., PRL 95 2005

**Isomerization** in terms of wave packet language:  
a ballistic passing of wave packet from excited state through conical intersection point (given as an “aperture”) to 13-cis ground state


\*V. Prokhorenko et al. Science 2006, 313: 1257

# Primary steps in bacteriorhodopsin photocycle: pump – probe kinetics of *all trans* → *13 cis* isomerization

Excitation with 20 fs laser pulse




2D-plot





## Samples:

- buffer NaCl + Phosph (pH = 6.5)
- OD in max. absorbance 0.8; flow cell 400  $\mu$ m
- room temperature, MA measuring conditions, cut-off filter (probe beam)
- light-adapted (before experiments and continuously during measurements)
- sonicated direct before measurements (for suppressing of scattering)

# Decay traces at different wavelengths



Scan in delay window 50 ps: shows some decay of *cis* – form during  $\sim$  few ps




Growth of *cis* – form occurs within  $\tau \sim 450$  fs

# Optimization experiment: enhancement of *cis* – yield using pulse shaping



Pump: 16 nJ, delay 20 ps after excitation; observation @ 630 nm (IF 10 nm)

Spectrum: controlled within 60 nm (540 – 600 nm), step 2 nm, 32 levels

Optimization process: 50 generations







Spectrum of optimal pulse




Spacing between main components:  
~ 7.1 nm (+/- 0.9 nm)  
→ ~ 210 cm<sup>-1</sup>

## Temporal structure of the optimal pulse: FROG data and Wigner plot





# Minimization experiment: suppressing *cis*-yield using shaped pulses





Anti-optimization process: 60 generations






Spectrum of anti-optimal pulse




## Temporal structure of the anti-optimal pulse



# Phase dependence: optimal pulse with - and without phase modulation





Spectra of pulses - identical



FROG of pulse with flat phase


⇒ Coherent Control....Quantum Coherence persists along reaction coordinate

# Phase Dependence: Reaction Dynamics




Phase dependence of the reaction branching ratio should be reflected in the reaction dynamics

# Pulse Shape Dependence of Molecular Dynamics



(A1,2) – without modulation; (B1,2) – with phase modulation; (C1,2) – with flipped phase modulation

- Analysis of pump-probe kinetics driven by different pulses






### EXAMPLE:



**Several traces at different wavelengths**  
 (note – actinic excitation energy all the same)

Blue – transform-limited  
 Red – optimal  
 Black – anti-optimal

## • Global Spectral Analysis




$$dA(t, \lambda) = IRF \otimes \sum_i A_i(\lambda) e^{-t/\tau_i}$$



### Basics:


population / isomerization kinetics are *sensitive* to phase information contained in light

# Coupling to Reaction Mode



→ Driving Large Amplitude Motion along Rxn Coordinate

# Mechanism: Insight from Theoretical Studies



S. Flores, V.  
Baptista, JPCB,  
2004, 6745  $\Rightarrow$   
pulse shape  
comprised of 2  
gaussians

**Figure 2.** Time-dependent probability of the system to be in the trans configuration of the  $S_0$  electronic state,  $P_{\text{trans}}^{(S_0)}(t)$  (upper panel), and the cis configuration of the  $S_1$  electronic state,  $P_{\text{cis}}^{(S_1)}(t)$  (lower panel), for relative pump pulse phases  $\theta = 0, 3\pi/2, \pi/2$ , and  $\pi$ .

TDSCF: Full quantum treatment (25) modes with empirical coupling to protein

$\Rightarrow$  Same excitation level as experiment: predicts 30% control

$\Rightarrow$  Time dependent reaction probability: material response is time variant viz bifurcation point in Conical Intersection

## CONCLUSIONS (CIRCA 2009)

---

- Trans-cis isomerization (branching ratio) of retinal molecule in bacteriorhodopsin **can be controlled** in weak field limit using tailored excitation pulses (40-50%)  
⇒ **control of a biological function**
- *Fundamental differences for weak field control in closed and open quantum systems*
- Optimal pulse displays very regular temporal- and spectral structure ⇒ coincides with driving torsional reaction mode modulating the conical intersection
  - central spectral components are modulated with period of ~ 150, 80, 45 fs

⇒ **Coherence is conserved through barrier crossing events in biological systems — and can be controlled/manipulated. “Proteins know how to surf”**

# EXTENSION TO STRONG FIELD: THE CHALLENGE

## Control of retinal isomerization in bacteriorhodopsin in the high-intensity regime


Andrei C. Florean<sup>a</sup>, David Cardoza<sup>b</sup>, James L. White<sup>b</sup>, J. K. Lanyi<sup>c</sup>, Roseanne J. Sension<sup>a,1</sup>, and Philip H. Bucksbaum<sup>b,1</sup>

<sup>a</sup>Department of Physics, University of Michigan, Ann Arbor, MI 48109; <sup>b</sup>PULSE Institute and Department of Physics, Stanford University, Stanford, CA 94305; and <sup>c</sup>School of Medicine, University of California, Irvine, CA 92697

Contributed by Philip H. Bucksbaum, May 20, 2009 (sent for review October 13, 2008)

A learning algorithm was used to manipulate optical pulse shapes and optimize retinal isomerization in bacteriorhodopsin, for excitation levels up to  $1.8 \times 10^{16}$  photons per square centimeter. Below 1/3 the maximum excitation level, the yield was not sensitive to pulse shape. Above this level the learning algorithm found that a Fourier-transform-limited (TL) pulse maximized the 13-cis population. For this optimal pulse the yield increases linearly with intensity well beyond the saturation of the first excited state. To understand these results we performed systematic searches varying the chirp and energy of the pump pulses while monitoring the isomerization yield. The results are interpreted including the influence of 1-photon and multiphoton transitions. The population dynamics in each intermediate conformation and the final branching ratio between the all-trans and 13-cis isomers are modified by changes in the pulse energy and duration.

coherent control | photoisomerization | ultrafast science



Coherent Control Absent in High Intensity Regime in contrast to all other systems ⇒ something different about biological systems >>> COMPLEXITY <<<<<<<<

Isomerization is more efficient from higher lying electronic states.

⇒ How can an upper level state, never accessed, be more efficient than evolutionary optimized state?.....contradicts weak field control results

# No Coherence in Control






Fig. 4. Probe absorption change vs. chirp of the pump pulse. (Upper) The  $J_{625}$  (blue diamonds) and  $K_{590}$  (red squares) signals versus linear chirp at 80 nJ pulse energy. (Lower)  $K_{590}$  signal versus linear chirp at 12 nJ pulse energy (green triangles). All signals are measured at 650 nm. The  $J_{625}$  signal is measured at 2 ps and the  $K_{590}$  signal is measured at 40 ps.

⇒Cis Formation Probed at 650 nm

Optimal Control Pulse is observed to be Transform Limited ⇒ No relative phase dependence, “control” only depends on peak power

Contradicts Weak Field Control Studies and generalized observation of increased control in strong fields

# Experiment Repeated: Chirp scans



← dA vs. chirp


← absorbed energy vs. chirp  
← Rxn Yield increases with  
Neg. Chirp

⇒ Phase Dependence is Observed and is Significant (16%)

⇒ Reproduced Results at Highest Intensity/Conditional Proof

⇒ Florean et al/wrong monitoring wavelength/not normalized to absorbed energy... rxn yield was not the observable

## Differential absorption spectra measured at 40 ps delay after excitation (sample OD ~ 1)




**Origin of observed spectral shift**  
⇒ ionization of bR and generation of solvated electrons

⇒ More than one photoproduct

# CONTROL STUDY — BUFFER ONLY


Chirp scan of very diluted sample (OD 0.2) measured @600 nm



Absorbed energy in pure buffer vs. chirp rate

Reproduces effect without protein → 10% of excitation absorbed due to  
**multiphoton absorption/ionization under NONRESONANT CONDITIONS >>>**  
Orders of Magnitude larger for RESONANT CONDITIONS of bR

# 1) Intrinsic isomerization control: wave packet acceleration



Parameter  $g = v^{3/2}$


$v$  – speed of wave packet going through the conical intersection “aperture” (i.e., chirp of pulse)

**Negatively-chirped pulses should increase isomerization efficiency**


Piryatinski et al., *PRL* 223001 (2005)

Negative chirp enhances motion to conical intersection...less time for scattering into unreactive modes

## 2) Control of Isomerization: High Intensity Regime (“Exact”)



Abe et al., J.Chem.Phys. 123, 144508 (2005)



- subpulses have a period of ~ 20 fs corresponding to a carbon backbone stretch of ~1600 cm<sup>-1</sup>  
Frozen two levels → does not include coupling to protein...15% for FC weighted wavepacket

**General Feature ⇒ optimum pulse is composed of subpulses timed to modes involved in reaction**

## CONCLUSIONS (CIRCA 2010)

---

Coherent Control demonstrated from weak field to strong field limits


⇒ *Fundamental differences for weak field control in closed and open quantum systems*

⇒ **Key Message: Protein Structure Reduces the Reaction Coordinate to a Few Labile Coordinates**

⇒ Coherent Control must be extended to Weak Field Limit to avoid multiphoton ionization/multiple reaction channels

⇒ *Coherence is conserved through barrier crossing events in biological systems — and can be controlled/manipulated. “Proteins know how to surf”*



***Nagging Question: How to rationalize degree of Coherent Control with 10 fs regime Quantum Decoherence of the Optically Induced Polarization?***



# Characterizing Quantum Coherence in Biological Systems ⇒ Coherent Multidimensional Spectroscopy


## Motivation:

- ⇒ Two-dimensional photon-echo electronic spectroscopy (2DPE) directly measures the homogeneous linewidth (pure dephasing, T2 contribution), couplings between states, and enables watching the state preparation evolve spectrally...more information on bR problem.
- ⇒ Anomolously long lived coherences have also been suggested to play a role in energy transport in photosynthetic systems...quantum or wave like transport...special role of the protein environment



Oberling, Strumpfer, Schulten, JPC, 2010

Panitchayangkoon et al. , PNAS, 2010



# Understanding 2D-PE spectra

## 1) Ensemble of identical molecules

$T = 0$  ("correlation spectrum")

$$S_{PE}(\omega_t, \omega_\tau) \propto \int_0^\infty dt e^{-2\text{Re}(g(t))} e^{i\omega_t t} \int_0^\infty d\tau \chi(t-\tau) e^{-2g^*(\tau)} e^{g^*(t+\tau)} e^{i\omega_\tau \tau}$$



correlations  
 $\rightarrow$  link between  $\omega_\tau, \omega_t$

$T \rightarrow \infty$  (no inhomogeneous broadening)

$$S_{PE}(\omega_t, \omega_\tau) \propto \sigma_a(\omega_\tau) \{ \sigma_a(\omega_t) + \sigma_f(\omega_t) + iKK[\sigma_a(\omega_t) + \sigma_f(\omega_t)] \}$$


$\rightarrow$  there is no link between variables!

Increasing  $T \rightarrow \infty$




## 2) Uncoupled molecules with different electronic transitions

Level diagram:




Density matrix (2 molecules):


$$\rho(t) = \begin{Bmatrix} \rho_{22} & 0 & \rho_{20} \cos(\omega_{20}t) \\ 0 & \rho_{11} & \rho_{10} \cos(\omega_{10}t) \\ \rho_{02} \sin(\omega_{20}t) & \rho_{01} \sin(\omega_{10}t) & \rho_{00} \end{Bmatrix}$$



## 3) Excitonically-coupled molecules (molecular aggregate)




$$\rho(t) = \begin{Bmatrix} \rho_{22} & \rho_{21} \cos(\omega_{21}t) & \rho_{20} \cos(\omega_{20}t) \\ \rho_{12} \sin(\omega_{12}t) & \rho_{11} & \rho_{10} \cos(\omega_{10}t) \\ \rho_{02} \sin(\omega_{20}t) & \rho_{01} \sin(\omega_{10}t) & \rho_{00} \end{Bmatrix}$$




# Example “TLS”: Rhodamine 101


$T = 0 \text{ fs}$



$T = 40 \text{ ps}$




# Model Dimer:





- Clearly resolved cross peaks – note amplitude is as expected from cross terms (e.g.  $\mu^2_A \mu^2_C$ )

# Quantum Beats/Homogeneous Lifetime




⇒ linewidth gives an electronic dephasing time of  $\sim 45$  fs...also in decay of photon echo ampl.



	AD	BD
$\omega$ (cm⁻¹)	1501	1550
$\tau$ (fs)	41.7	49.2


⇒ The antidiagonal line width and off diagonal components are causally related (FT)...long lived quantum beats are vibrational (Jonas et al – vibrational coherences enhance ET)

# Bacteriorhodopsin



negative feature growing  
in near cis max


**T = 0 fs**



- clear vibronic structure at HOOP frequency
- negative Kerr effect
- anti-diagonal linewidth:  $936 \text{ cm}^{-1}$ , results in a dephasing time of 11 fs  
**(upper bound)**

# $T = 10 \text{ fs}$

negative feature not due  
to ESA




- negative feature **grows**
- located (spectrally) at the *cis* absorption max!
- correlated with vibronic bleach feature
- vibronic band shows off-diagonal **coupling**

# Temporal dynamics


Effect of pumping about the vibrational shoulder at  $18500\text{cm}^{-1}$ : **clear oscillatory dynamics of the cis-like feature**

Effect of pumping the linear absorption maximum at  $17500\text{cm}^{-1}$ : **vibrational cross-peak**



# Fit results

$$\sum_{i=1}^4 A_i e^{-t/a_i} \sin \left( \frac{2\pi t}{b_i} + c_i \right) + d$$



Frequency, cm <sup>-1</sup>	Phase, rad	Amplitude, a.u.	Decay, fs	Mode
823	1.87	0.838	54	HOOP
210	2.73	0.731	40	Torsion
1560	1.09	0.453	53	C=C stretch
1106	2.20	0.935	42	C-C stretch
986	1.95	0.772	29	HOOP

- ⇒ **Very strong coupling between trans and cis electronic surfaces by the very modes directing the reaction coordinate**
- ⇒ **HIGHLY DIRECTED**

## CONCLUSIONS

---

Coherent Control demonstrated from weak field to strong field limits

⇒ *Fundamental differences for weak field control in closed and open quantum systems*

⇒ **Key Message:** Protein Structure Reduces the Reaction Coordinate to a Few Labile Coordinates

⇒ Coherent Control must be extended to Weak Field Limit to avoid multiphoton ionization/multiple reaction channels

⇒ *Coherence is conserved through barrier crossing events in biological systems — and can be controlled/manipulated.*  
“Proteins know how to surf”

***Vibrational Coupling/Coherences exploited for optimizing reaction coordinates/functions in biological systems***