
Isotropic Tensors, NS Equations, and Gauss' Theorem

0 Course details

- Course hours and location: Perlman 404 (or Zoom). Lectures on Tuesdays @ 11:15 and tutorial sessions on Thursdays @ 9:15.
 - Lecture zoom link:
<https://weizmann.zoom.us/j/94852313362?pwd=S0dBaXhOSVJudlpHK0N5UXl3eTZpUT09>
 - Tutorial session zoom link:
<https://weizmann.zoom.us/j/99636463168?pwd=VEYvRGVOWWcyOEpZYyt4NHN0bHVUQT09>
- Course website:
<https://www.weizmann.ac.il/chembiophys/bouchbinder/courses-0>.
- Course material will be uploaded to the website weekly (lecture notes, tutorial notes, exercises, and solutions to previous exercises). Please get updated via the course website above.
- Teaching Assistants: Avraham Moriel and Yuri Lubomirsky (our office @ Perlman 720). If needed, contact us via e-mail to schedule a meeting in advance.
- E-mail for homework submission, questions, suggestions and complaints:
groupbouchbinder@gmail.com
- Please feel free (and you are encouraged) to ask anything via email. We will use part of the tutorials to go over important questions and topics raised.
- Homework submission:
 - All homework submission is via the e-mail address above.
 - Please save your homework files with the following name:
NECP_HW#-(FirstName)-(LastName).pdf.
For example: NECP_HW3-Avraham_Moriel.pdf.
Please use the same format for the e-mail title (e.g. NECP_HW3-Avraham_Moriel).
 - We **strongly recommend** submitting computer-typed homeworks. This (1) helps us read and understand your solutions, and (2) will get you to experience scientific writing in its most basic form.
- We are new to the business of hybrid teaching — please share with us your past experience!

The purpose of today's TA session is to mess a bit with tensors and indices, which are a necessary tool for continuum theories and in particular for Solid Mechanics. We'll see some simple examples and try to become comfortable with these mathematical tools. If time permits we'll see a generalization of Gauss's Theorem to tensors.

1 Isotropic tensors

Eran defined tensors as linear operators transforming n into m vectors. One can define a tensor as an object that under orthogonal (unitary) coordinate transformations (i.e. rotations) transforms as

$$A_{i_1 i_2 \dots i_k} = Q_{i_1 j_1} Q_{i_2 j_2} \dots Q_{i_k j_k} A_{j_1 j_2 \dots j_k} , \quad (1)$$

where the Q 's represent the orthogonal transformation from coordinates j to coordinates i . We will not bother with the distinction between covariant and contravariant degrees of freedom (though they are crucial in other fields of physics like general relativity).

A tensor is called *isotropic* if its coordinate representation is independent under coordinate rotation. Let's look at all the possible forms of isotropic tensors of low ranks.

1.0 0th rank tensors

A 0th rank tensor, a.k.a a scalar, does not change under rotations, therefore all scalars are isotropic (surprise!).

1.1 1st rank tensors

A vector \vec{v} is isotropic if for every rotation matrix R_{ij} we have

$$R_{ij} v_j = v_i . \quad (2)$$

You can easily show that this condition is satisfied for arbitrary \mathbf{R} only if $\vec{v} = 0$. So the zero vector is the only isotropic vector (surprise #2!!).

1.2 2nd rank tensors

Let's hope we're gonna get something a bit more interesting. A matrix \mathbf{A} is isotropic if for every rotation matrix \mathbf{R} we have $A_{ij} = R_{ik} R_{jl} A_{kl}$, or in matrix notation:

$$\mathbf{R} \mathbf{A} \mathbf{R}^T = \mathbf{A} . \quad (3)$$

Let's choose a specific rotation matrix, say a rotation of angle α around \hat{z} ,

$$\mathbf{R}^z(\alpha) \equiv \begin{pmatrix} \cos \alpha & \sin \alpha & 0 \\ -\sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{pmatrix} . \quad (4)$$

The invariance equation now takes the form

$$\mathbf{A}(0) = \mathbf{A}(\alpha) \equiv \mathbf{R}^z(\alpha) \mathbf{A} \mathbf{R}^z(\alpha)^T = \begin{pmatrix} \cos \alpha & \sin \alpha & 0 \\ -\sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{pmatrix} \mathbf{A} \begin{pmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{pmatrix} . \quad (5)$$

This is a complicated equation, with cosines and sines all mixed up in a very unpleasant manner. Luckily, we can find an equivalent condition that is significantly simpler. Differentiating with respect to α and plugging $\alpha = 0$ gives

$$0 = \frac{\partial \mathbf{A}(0)}{\partial \alpha} = \left. \frac{\partial \mathbf{A}(\alpha)}{\partial \alpha} \right|_{\alpha=0} = \left. \frac{\partial \mathbf{R}^z(\alpha)}{\partial \alpha} \right|_{\alpha=0} \mathbf{A} \mathbf{R}^z(0) + \mathbf{R}^z(0) \mathbf{A} \left. \frac{\partial \mathbf{R}^z(\alpha)^T}{\partial \alpha} \right|_{\alpha=0}, \quad (6)$$

but since $\mathbf{R}^z(0)$ is the identity matrix, this reduces to the simple equation

$$\underbrace{\begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}}_{\equiv \mathbf{L}^z} \mathbf{A} + \mathbf{A} \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} = 0. \quad (7)$$

$\mathbf{L}^z = \partial_\alpha \mathbf{R}|_{\alpha=0}$ is sometimes called “the generator of rotations around the z axis”, because $\mathbf{R}^z(\alpha) = e^{\alpha \mathbf{L}^z}$. We see that equation $\mathbf{A}(0) = \mathbf{A}(\alpha)$ is equivalent to the much easier equation (notice the sign change)

$$\mathbf{A}(0) = \mathbf{A}(\alpha) \iff [\mathbf{A}, \mathbf{L}^z] = 0. \quad (8)$$

Explicitly calculating $[\mathbf{A}, \mathbf{L}^z]$ gives

$$[\mathbf{A}, \mathbf{L}^z] = \begin{pmatrix} -A_{12} - A_{21} & A_{11} - A_{22} & -A_{23} \\ A_{11} - A_{22} & A_{12} + A_{21} & A_{13} \\ -A_{32} & A_{31} & 0 \end{pmatrix}. \quad (9)$$

We see that commutation with \mathbf{L}^z requires (a) $A_{13} = A_{31} = A_{23} = A_{32} = 0$ and (b) $A_{11} = A_{22}$. Obviously, the choice of \hat{z} is arbitrary and isotropy means that \mathbf{A} should also commute with \mathbf{L}^x and \mathbf{L}^y . If we repeat the above procedure for the other \mathbf{L} 's, the analog of (a) will be that all off-diagonal elements must vanish, and the analog of (b) will be that all diagonal elements must be equal. That is,

$$A_{ij} \propto \delta_{ij}. \quad (10)$$

I stress that this is true only in dimensions ≥ 3 . In the HW you'll see that in 2D there are isotropic tensors that are not proportional to the identity (can you already see how the above argument fails in 2D?).

1.3 3rd rank tensors

Here we can use the same trick. A 3rd rank tensor \mathbf{A} is isotropic iff for every rotation matrix R_{ij} we have

$$R_{i\alpha} R_{j\beta} R_{k\gamma} A_{\alpha\beta\gamma} = A_{ijk}. \quad (11)$$

You can imagine the mess that comes out of this if you plug in a real rotation matrix with sines and cosines and whatnot, and then start using trig identities. Phew, no thanks!

So like before, we choose $\mathbf{R} = \mathbf{R}^z(\alpha)$, differentiate, and set $\alpha = 0$. This gives

$$\begin{aligned} 0 &= \left(L_{i\alpha}^z \delta_{j\beta} \delta_{k\gamma} + \delta_{i\alpha} L_{j\beta}^z \delta_{k\gamma} + \delta_{i\alpha} \delta_{j\beta} L_{k\gamma}^z \right) A_{\alpha\beta\gamma} \\ &= L_{i\alpha}^z A_{\alpha j k} + L_{j\beta}^z A_{i \beta k} + L_{k\gamma}^z A_{i j \gamma}. \end{aligned} \quad (12)$$

To see what kind of equation we got, let's choose $i = 1, j = 3, k = 3$. Since the only non-zero elements of \mathbf{L}^z are L_{12}^z and L_{21}^z , we get

$$0 = L_{1\alpha}^z A_{\alpha 33} + L_{3\beta}^z A_{1\beta 3} + L_{3\gamma}^z A_{13\gamma} = A_{233} . \quad (13)$$

Similarly, by choosing different combinations of i, j, k and/or different \mathbf{L} 's, you get that $A_{ijk} = 0$ whenever i, j, k are not all different, that is, if (ijk) is not a permutation of (123) .

Using this knowledge, we can choose now $i = 1, j = 1, k = 3$, and we get

$$A_{113} = 0 = L_{1\alpha}^z A_{\alpha 13} + L_{1\beta}^z A_{1\beta 3} + L_{3\gamma}^z A_{11\gamma} = A_{213} + A_{123} ,$$

or put differently, $A_{213} = -A_{123}$. Similarly, we can show that every time we flip two indices we get a minus sign. Therefore, we conclude that the only isotropic 3rd rank tensor is equal, up to a multiplicative constant, to \mathcal{E} ,

$$\mathcal{E}_{ijk} = \begin{cases} 0 & (ijk) \text{ is not a permutation of } (123) \\ \text{sign of permutation} & \text{otherwise} \end{cases} . \quad (14)$$

As you probably know, \mathcal{E} is called the Levi-Civita completely anti-symmetric tensor¹.

1.4 4th rank tensors

Seriously? No. We're not going to redo the algebra. But can we guess the form of some isotropic 4th rank tensors? We can easily build them from lower rank isotropic tensors. Here are a few examples that come to mind:

$$A_{ijkl} = \delta_{ij} \delta_{kl} , \quad (15)$$

$$A_{ijkl} = \delta_{il} \delta_{jk} , \quad (16)$$

$$A_{ijkl} = \delta_{ik} \delta_{jl} , \quad (17)$$

$$A_{ijkl} = \mathcal{E}_{ij\alpha} \mathcal{E}_{\alpha kl} . \quad (18)$$

We did a really good job there, because it turns out that these are the only options. In fact, this list is even redundant, because each of the lines can be written as a linear combination of the other three (can you find it?). You may want to prove at home that there really are no other options - it's a nice exercise that can be easily automatized on *Mathematica*, and we're going to use this result in the course.

A generic approach to produce isotropic tensors is based on group theory, and specifically the Clebsch–Gordan decomposition of tensor product (I suggest "Group Theory in a Nutshell for Physicists" by A. Zee).

¹ This is true only in flat spaces. Those of you familiar with differential geometry might insist on calling it a "Tensor density". Since we are (thankfully) only considering flat space here, we'll disregard this subtlety.

2 Navier-Stokes equation

We are now going to use the heavy arsenal developed above, and derive the Navier-Stokes (NS) equation solely from symmetry considerations. We want to find a dynamical equation for $\partial_t \vec{v}$ as a function of \vec{v} and its spatial derivatives. We take a perturbative approach, and expand $\partial_t v$ to second order in \vec{v} and in its gradients:

$$\partial_t v_i = A_{ij} v_j + B_{ijk} \partial_j v_k + D_{ijkl} v_j \partial_k v_l + E_{ijkl} \partial_j \partial_k v_l + F_{ijk} v_j v_k + G_{jkl} \partial_j v_i \partial_k v_l . \quad (19)$$

Since \vec{v} is a physical quantity (specifically, a 1st rank tensor), the dynamical equation for $\partial_t \vec{v}$ should be covariant under symmetries of the physical system in question. We'll see what these symmetries impose on the form of the various tensors $\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}, \mathbf{F}, \mathbf{G}$.

We begin with a Galilean transformation:

$$y_i = x_i - c_i t , \quad (20)$$

$$\tau = t . \quad (21)$$

Under this transformation, the velocity field now takes the form $\vec{w} = \vec{v} - \vec{c}$. Also, by the chain rule:

$$\partial_{x_i} = \frac{\partial y_j}{\partial x_i} \partial_{y_j} + \frac{\partial \tau}{\partial x_i} \partial_\tau = \partial_{y_i} , \quad (22)$$

$$\partial_t = \frac{\partial \tau}{\partial t} \partial_\tau + \frac{\partial y_j}{\partial t} \partial_{y_j} = \partial_\tau - c_j \partial_{y_j} . \quad (23)$$

Applying this to Eq. (19) gives

$$\begin{aligned} \partial_t w_i - c_j \partial_j w_i &= A_{ij} (w_j + c_j) + B_{ijk} \partial_j w_k + D_{ijkl} (w_j + c_j) \partial_k w_l \\ &\quad + E_{ijkl} \partial_j \partial_k w_l + F_{ijk} (w_j + c_j) (w_k + c_k) + G_{jkl} \partial_j w_i \partial_k w_l . \end{aligned} \quad (24)$$

If we want the NS equation to be covariant, we need to impose that Eq. (24) will be equal, term by term, to Eq. (19), i.e.

$$A_{ij} c_j = 0 , \quad (25)$$

$$-c_j \partial_j w_i = D_{ijkl} c_j \partial_k w_l , \quad (26)$$

$$F_{ijk} (w_j + c_j) (w_k + c_k) = F_{ijk} w_j w_k . \quad (27)$$

All these should hold for arbitrary \vec{c}, \vec{w} . The first constraint clearly means $\mathbf{A} = 0$. For the third one, choose for example $\vec{w} = -\vec{c}$, and get that $F_{ijk} w_j w_k = 0$ for arbitrary \vec{w} . Note that this is exactly the last term in Eq. (19), so we see that it vanishes identically. The constraint Eq. (26) may be written as

$$-\delta_{il} \delta_{jk} c_j \partial_k w_l = D_{ijkl} c_j \partial_k w_l .$$

Since \vec{c}, \vec{w} are arbitrary, $D_{ijkl} = -\delta_{il} \delta_{kj}$, and Eq. (19) can be written as

$$\partial_t v_i + v_j \partial_j v_i = B_{ijk} \partial_j v_k + E_{ijkl} \partial_j \partial_k v_l + G_{jkl} \partial_j v_i \partial_k v_l . \quad (28)$$

You have to admit that this is a very big improvement...

Now let's look at rotations $y_j = R_{ij}x_j$. Demanding Eq. (28) to be invariant means that the tensors \mathbf{B} , \mathbf{E} and \mathbf{G} are *isotropic*.

We've just seen that the only 3rd rank isotropic tensor is the Levi-Civita tensor, so the \mathbf{B} and \mathbf{G} terms are proportional to $\vec{\nabla} \times \vec{v}$ and thus are forbidden by reflection symmetry. It's too bad that we know already that the \mathbf{A} and \mathbf{F} terms are gone, because they would also be forbidden by rotational symmetry. For example, the \mathbf{F} term must be proportional to $\vec{v} \times \vec{v}$ and therefore vanishes identically (note that we didn't show that $\mathbf{F} = 0$, but only that it gives zero when it acts on the same vector in its two slots).

As for \mathbf{E} , we know that we have exactly three choices, given in Eqs. (15),(16),(17). These give, respectively,

$$\delta_{ij} \delta_{kl} \partial_j \partial_k v_l = \partial_i \partial_j v_j = \vec{\nabla} (\nabla \cdot \vec{v}) = \text{grad} (\text{div} \vec{v}) , \quad (29)$$

$$\delta_{il} \delta_{jk} \partial_j \partial_k v_l = \partial_j \partial_j v_i = \nabla^2 \vec{v} = \text{div} (\text{grad} \vec{v}) , \quad (30)$$

$$\delta_{ik} \delta_{jl} \partial_j \partial_k v_l = \partial_i \partial_j v_j = \text{same as Eq. (29)} , \quad (31)$$

so the third option is redundant. Note that if we wanted to use Eq. (18) we'd get

$$\mathcal{E}_{ij\alpha} \mathcal{E}_{\alpha kl} \partial_j \partial_k v_l = \mathcal{E}_{ij\alpha} \partial_j \left(\vec{\nabla} \times \vec{v} \right)_\alpha = \vec{\nabla} \times \left(\vec{\nabla} \times \vec{v} \right) ,$$

which is also redundant because of the vector calculus identity which you all know by heart: $\nabla \times (\nabla \times \mathbf{A}) = \nabla(\nabla \cdot \mathbf{A}) - \nabla^2 \mathbf{A}$.

To sum up, we see that the only form of $\partial_t \vec{v}$ which is covariant under rotations, reflections and Galilean symmetries is

$$\left(\partial_t + \vec{v} \cdot \vec{\nabla} \right) \vec{v} = \eta \nabla^2 \vec{v} + \mu \vec{\nabla} (\nabla \cdot \vec{v}) , \quad (32)$$

where η and μ are two scalars. In incompressible flows, $(\nabla \cdot \vec{v})$, there's only one η , as the μ term vanishes.

Lastly, note that there's another term that clearly does not violate any symmetries: $\vec{\nabla} P$ where P is some scalar function.

2.1 A historical note about the power of symmetries in continuum theories

Euler's equation $(\partial_t + v_j \partial_j) v_i = \partial_i P$, regarding inviscid incompressible flows, was derived sometime around 1750. It took the scientific community almost 80 years (!) to understand how to incorporate viscosity into the business. Mind you, some of the greatest minds of the time were devoted to the problem, including Cauchy, Poisson, d'Alembert, Bernoulli, and of course, Navier and Stokes. Not exactly Elitzur Ra'anana, if you see what I mean. So what took them so long?

The answer, very very roughly, is that they tried to model viscosity on a molecular level: to understand the dissipation mechanisms, stress-transfer mechanisms, and what-not. One of the great strengths of continuum theory is that measly insignificant mortals like us were able to do here in 45 minutes a derivation that the primordial gods needed 80 years to do. Moreover, we did that *without caring even the slightest bit about the underlying physics*.

In fact, this is the crux of the matter – the use of symmetries allows us to state very powerful statements about the functional form of the viscosity term, without having to deal with the microscopic mechanisms. It allows us to develop a predictive theory, where all the “microscopics” are lumped into a small number of parameters (in our case - μ and η), which of course must be determined experimentally.

The down side is that we can not say anything quantitative about the parameters. From our theory we can not give even an order-of-magnitude estimation of η or μ , let alone their dependence on the fluid’s properties (although thermodynamics easily tells us that they are positive).