

- (Ch. 3, EOC ex. 3) Let $\psi(x, 0) = e^{-\alpha x^2}$. Find an expression for $\psi(x, t)$ by using the Fourier integral theorem. Show that $\psi(x, t)$ can be written in the form $\psi(x, t) = e^{-\alpha_t x^2 + \gamma_t}$.
- (Ch. 3, EOC ex. 4) An alternative approach to the previous problem is: guess that $\psi(x, t) = e^{-\alpha_t x^2 + \gamma_t}$. Substitute this form into the time dependent Schrodinger equation and, by matching equations in powers of x find expressions for α_t and γ_t . Show that these expressions are equal to those derived in the previous problem.
- (Ch. 3, EOC ex. 5) Take the more general trial form $\psi(x, t) = e^{-\alpha_t(x-x_t)^2 + \frac{i}{\hbar}p_t(x-x_t) + \frac{i}{\hbar}\gamma_t}$, where x_t and p_t are real. Substitute this form into the TDSE and obtain expressions for α_t , x_t , p_t , and γ_t .
- (Ch. 3 EOC ex. 6) Find $\overline{x}(t)$ and $\overline{\Delta x^2}(t)$, $\overline{p}(t)$ and $\overline{\Delta p^2}(t)$ for exercise 2 above. Show that the uncertainty principle is satisfied:

$$\Delta x(t)\Delta p(t) \geq \frac{\hbar}{2}. \quad (4.1)$$

- (Ch. 3, EOC ex. 8) Take the form for the gaussian wavepacket, $\psi(x, t) = \exp(-\alpha_t(x-x_t)^2 + \frac{i}{\hbar}p_t(x-x_t) + \frac{i}{\hbar}\gamma_t)$, and substitute it into the Time Dependent Schrödinger Equation for the harmonic oscillator potential, $V(x) = \frac{1}{2}m\omega^2 x^2$. Obtain expressions for the parameters α_t , x_t , p_t , and γ_t . [Hint: to simplify the algebra, write $\frac{1}{2}m\omega^2 x^2 = \frac{1}{2}m\omega^2[(x-x_t) + x_t]^2$. To solve the differential equation for α_t and γ_t it is convenient to introduce the substitution $\alpha_t = \frac{a\dot{z}}{z}$, where a is a constant which gives the harmonic equation $\ddot{z} = -\omega z$.] What happens when $\alpha_0 = \frac{m\omega}{2\hbar}$? In this case the wavepacket is sometimes called a 'coherent state', a confusing but widely used label.
- (Ch. 3, EOC ex. 1) Optional: Show that the product of two gaussians is a third gaussian centered between the first two.
- (Ch. 3, EOC ex. 2) Optional: Find an expression for the overlap integral of two complex gaussians of the form $\psi(x, t) = e^{-\alpha_t(x-x_t)^2 + \frac{i}{\hbar}p_t(x-x_t) + \frac{i}{\hbar}\gamma_t}$, the first with center at x', p' , and the second with center at x'', p'' . Note that this expression depends only on the differences $|p' - p''|$ and $|x' - x''|$. Interpret the exponent in terms of the phase space distance between the centers of the two gaussians by finding a quantitative relationship between the two.
- (Ch. 3, EOC ex. 7) Optional: Take the form for the gaussian wavepacket, $\psi(x, t) = \exp(-\alpha_t(x-x_t)^2 + \frac{i}{\hbar}p_t(x-x_t) + \frac{i}{\hbar}\gamma_t)$, and substitute it into the Time Dependent Schrödinger Equation for the linear potential, $V(x) = -kx$. Obtain expressions for the parameters α_t , x_t , p_t , and γ_t .
- (Ch. 3, EOC ex. 9) Optional: Consider the forced harmonic oscillator, *i.e.*, $V(x) = 1/2m\omega^2 x^2 - F(t)x$.
 - Show that the wavepacket given by $\psi(x, t) = \exp(-\alpha_t(x-x_t)^2 + \frac{i}{\hbar}p_t(x-x_t) + \frac{i}{\hbar}\gamma_t)$ is a solution to this problem. Substitute this form into the Time Dependent Schrödinger Equation and obtain expressions for x_t , p_t , α_t , and γ_t .
 - A convenient way of solving the differential equations for x_t and p_t is to define a complex quantity $z(t) = p_t + im\omega x_t$. Then $z(t) = e^{i\omega t} \int_0^t F(t')e^{-i\omega t'} dt' + e^{i\omega t} z(0)$. Differentiate $z(t)$ with respect to time and show that the resulting equation for $\dot{z}(t)$ is consistent with Hamilton's equations of motion for x_t and p_t .
 - Solve for $z(t)$ for
 - $F(t) = \sin(\omega t)$
 - $F(t) = A$

iii. $F(t) = At$.

Draw pictures in the complex plane of $z(t)$ for all these cases.

- d. Show that in general, if $t(dF/dt) \ll F$, then $z(t) = iF(t)/\omega$ (taking $x_0 = p_0 = 0$). *Hint: Integrate by parts.* This is the limit of adiabatic forcing, i.e. if the change in the Hamiltonian is slow enough an initial eigenstate will stay an eigenstate for all time.