


The Non Equilibrium Steady State of Sparse Systems with Non Trivial Topology

Daniel Hurowitz

Department of Physics, Ben Gurion University, Beer Sheva

$$\mathcal{H}_{\text{total}} = \text{diag}\{E_n\} - f(t)\{V_{nm}\} + F \cdot \{W_{nm}\} + \mathcal{H}_{\text{Bath}}$$

- ▶ Doron Cohen (BGU) [1,2]
- ▶ Saar Rahav (Technion)[2]





1. D. Hurowitz and D. Cohen, Europhysics Letters 93, 60002 (2011)
2. D. Hurowitz, S. Rahav and D. Cohen, Europhysics Letters 98, 20002 (2012).

“Sparsity”

$$\mathcal{H}_{\text{total}} = \text{diag}\{E_n\} - f(t)\{V_{nm}\} + F \cdot \{W_{nm}\} + \mathcal{H}_{\text{Bath}}$$

Histogram of couplings





$$g_n = |V_{n,n+1}|^2 = \text{couplings}$$

“sparsity” = log wide distribution of couplings

Current vs. driving

Driving \rightsquigarrow Stochastic Motive Force \rightsquigarrow Current

Regimes: LRT regime, Sinai regime, Saturation regime



$$I \sim \frac{1}{N} \bar{w} \exp \left[-\frac{\mathcal{E}_n}{2} \right] 2 \sinh \left(\frac{\mathcal{E}_0}{2} \right)$$

Extent of the “Sinai regime” is determined by width of distribution of rates

Master equation description of dynamics

$$\mathcal{H}_{\text{total}} = \text{diag}\{E_n\} - f(t)\{V_{nm}\} + F \cdot \{W_{nm}\} + \mathcal{H}_{\text{Bath}}$$

Quantum master equation for the reduced probability matrix:

$$\frac{d\rho}{dt} = -i[\mathcal{H}_0, \rho] - \frac{\epsilon^2}{2}[V, [V, \rho]] + \mathcal{W}^\beta \rho \equiv \mathcal{W}\rho$$

Stochastic rate equation:

$$\frac{dp_n}{dt} = \sum_m w_{nm} p_m - w_{mn} p_n$$

The transition rates:

$$w_{nm} = w_{nm}^\epsilon + w_{nm}^\beta$$

$$w_{nm}^\epsilon = w_{mn}^\epsilon = g_{nm} \epsilon^2$$

Steady state equation:

$$\dot{\rho} = \mathcal{W}\rho = 0$$

$$\frac{w_{nm}^\beta}{w_{mn}^\beta} = \exp\left[-\frac{E_n - E_m}{T_B}\right]$$

The Stochastic Motive Force (SMF)

If we had only a bath

$$\frac{w_{nm}}{w_{mn}} = \exp \left[-\frac{E_n - E_m}{T_B} \right]$$

We define a “field”


$$\mathcal{E}(x) \equiv \ln \left[\frac{w_{nm}}{w_{mn}} \right]$$

and “potentials”

$$\mathcal{E}(x_1 \rightsquigarrow x_2) = \int_{x_1}^{x_2} \mathcal{E}(x) dx \quad \text{[potential variation]}$$

$$\mathcal{E}_{\cap} \equiv \text{maximum} \left\{ |\mathcal{E}(x_1 \rightsquigarrow x_2)| \right\} \quad \text{[activation barrier]}$$

$$\mathcal{E}_{\circlearrowleft} \equiv \oint \mathcal{E}(x) dx \quad \text{if no driving} = 0 \quad \text{[SMF]}$$





With driving, $\mathcal{E}_{\circlearrowleft} \neq 0$. This means $\prod_n w_{n,n+1} \neq \prod_n w_{n+1,n}$.

Current vs. driving

Driving \rightsquigarrow Stochastic Motive Force \rightsquigarrow Current

Regimes: LRT regime, Sinai regime, Saturation regime



$$I \sim \frac{1}{N} \bar{w} \exp \left[-\frac{\mathcal{E}_n}{2} \right] 2 \sinh \left(\frac{\mathcal{E}_0}{2} \right)$$

Extent of the “Sinai regime” is determined by width of distribution of rates

Emergence of the “Sinai regime”

Sinai [1982]: Transport in a chain with random transition rates.

Assume transition rates are uncorrelated.

↪ build up of a potential barrier $\mathcal{E}_n \propto \sqrt{N}$.

↪ exponentially small current.

But... we have telescopic correlations: $\mathcal{E}_{n,n+1} \sim \Delta_n \equiv (E_n - E_{n+1})$

Yet... we have sparsely distributed couplings: $w_{n,n+1}^\epsilon = g_n \epsilon^2$


$$\mathcal{E}_0 \approx - \sum_n \left[\frac{1}{1 + g_n \epsilon^2} \right] \frac{\Delta_n}{T_B} \sim \frac{1}{T_B} \begin{cases} \epsilon^2, & \epsilon^2 < 1/g_{\max} \\ 1/\epsilon^2, & \epsilon^2 > 1/g_{\min} \\ [\pm] \sqrt{N} \Delta, & \text{otherwise} \end{cases}$$

Build up may occur if g_n are from a **log-wide** distribution.

$$I \sim \frac{1}{N} \bar{w} \exp \left[-\frac{\mathcal{E}_0}{2} \right] 2 \sinh \left(\frac{\mathcal{E}_0}{2} \right)$$

Beyond fluctuation dissipation phenomenology: Topological term in EAR formula

$$\begin{aligned}\dot{Q} &= \sum_n \left[w_n^\beta p_n - w_n^\beta p_{n-1} \right] \Delta_n \\ &\approx \left[\frac{D_B}{T_B} - \frac{D_B}{T(0)} \right] + T_B \mathcal{E} \circ I \\ &\approx \frac{D_B}{T_B} \left[(g_n \epsilon^2) - (g_n \epsilon^2)^2 + \text{Var}(g) \epsilon^4 \right]\end{aligned}$$



The EAR is correlated with the current.

The quantum mechanical steady state


Stochastic

$$\frac{dp_n}{dt} = \sum_m w_{nm} p_m - w_{mn} p_n$$

$$I_{n \rightarrow m} = w_{mn} p_n - w_{nm} p_m \equiv \text{tr}(\hat{I}_{n \rightarrow m} \rho)$$

$$\hat{I}_{n \rightarrow m}^\epsilon = |n\rangle w_{mn}^\epsilon \langle n| - |m\rangle w_{nm}^\epsilon \langle m|$$

$$\hat{I}_{n \rightarrow m}^\beta = |n\rangle w_{mn}^\beta \langle n| - |m\rangle w_{nm}^\beta \langle m|$$




Quantum

$$\frac{d\rho}{dt} = -i[\mathcal{H}_0, \rho] - \frac{\epsilon^2}{2}[V, [V, \rho]] + \mathcal{W}^\beta \rho$$

$$\hat{I}_{n \rightarrow m}^\epsilon = i\epsilon^2 [\hat{J}^{nm}, \hat{V}]$$

$$\hat{J}^{nm} = i(|m\rangle V_{mn} \langle n| - |n\rangle V_{nm} \langle m|)$$



Summary of main results

1. Due to the **sparsity** of the perturbation matrix, the NESS is of glassy nature [1].
2. An extension of the Fluctuation-Dissipation phenomenology has been proposed [1].
3. A log-wide distribution of couplings is required in order to have a **Sinai regime**.
4. The **topological term** in the EAR is correlated with the current but sub-linear in driving intensity.
5. Novel saturation effect in the quantum model.
6. The quantum **current operator** in the reduced description includes off diagonal elements of the probability matrix.

[1] D. Hurowitz and D. Cohen, Europhysics Letters 93, 60002 (2011).

References and Acknowledgements

1. D. Hurowitz and D. Cohen, Europhysics Letters 93, 60002 (2011).
 2. D. Hurowitz, S. Rahav and D. Cohen, Europhysics Letters 98, 20002 (2012).
- ▶ **Sparsity**: Austin, Wilkinson, Prosen, Robnik, Alhassid, Levine, Fyodorov, Chubykalo, Izrailev, Casati
 - ▶ **Energy absorption by sparse systems**: Cohen, Kottos, Schanz, Wilkinson, Mehlig
 - ▶ **Network theory**: Schnakenberg, Zia, Hill
 - ▶ **Sinai physics**: Sinai, Derrida, Pomeau, Burlatsky, Oshanin, Mogutov, Moreau, Bouchard

Acknowledgement: Bernard Derrida