

Constraints on Exotic Dipole-Dipole Couplings between Electrons at the Micrometer Scale

Shlomi Kotler,¹ Roee Ozeri,² and Derek F. Jackson Kimball³

¹*National Institute of Standards and Technology, 325 Broadway Street, Boulder, Colorado 80305, USA*

²*Department of Physics of Complex Systems, Weizmann Institute of Science, P.O. Box 26, Rehovot 76100, Israel*

³*Department of Physics, California State University, East Bay, Hayward, California 94542-3084, USA*

(Received 30 January 2015; published 17 August 2015)

New constraints on exotic dipole-dipole interactions between electrons at the micrometer scale are established, based on a recent measurement of the magnetic interaction between two trapped $^{88}\text{Sr}^+$ ions. For light bosons (mass ≤ 0.1 eV) we obtain a 90% confidence interval for an axial-vector-mediated interaction strength of $|g_A^e g_A^e / 4\pi\hbar c| \leq 1.2 \times 10^{-17}$. Assuming *CPT* invariance, this constraint is compared to that on anomalous electron-positron interactions, derived from positronium hyperfine spectroscopy. We find that the electron-electron constraint is 6 orders of magnitude more stringent than the electron-positron counterpart. Bounds on pseudoscalar-mediated interaction as well as on torsion gravity are also derived and compared with previous work performed at different length scales. Our constraints benefit from the high controllability of the experimental system which contained only two trapped particles. It therefore suggests a useful new platform for exotic particle searches, complementing other experimental efforts.

DOI: 10.1103/PhysRevLett.115.081801

PACS numbers: 14.80.Va, 03.67.Pp, 14.60.Cd, 42.50.Dv

Extensions of the standard model which predict new particles may involve modifications to the known spin-spin interaction between fermions [1]. Examples include pseudoscalar fields (such as the axion [2]) which naturally emerge from theories with spontaneously broken symmetries [3–5], and axial-vector fields such as paraphotons [6] and extra Z bosons [1,7], which appear in new gauge theories. Both pseudoscalar and axial-vector fields are candidates to explain dark matter [8], dark energy [9,10], mysteries surrounding *CP* violation [2], the hierarchy problem [11], and are a generic prediction of string theories [12].

Typically, to constrain the effect of a new particle of mass m it is favorable to perform an experimental investigation at a length scale $\lambda \equiv \hbar/mc$, the reduced Compton wavelength associated with the particle. Here, \hbar is Planck's constant divided by 2π and c is the speed of light. At the macroscopic scale, there have been a number of searches for exotic dipole-dipole interactions between electrons [13–19], ranging from distance scales of a few cm to the radius of the Earth. The large scales involved contributed to the exquisite sensitivity of these endeavors, allowing for signal averaging over a large number of spins. Such an approach, however, cannot be implemented for scales significantly smaller than a millimeter. Attempting to extrapolate these bounds to the micrometer scale also fails, due to the interactions' distance scaling [20].

The magnetic dipole-dipole interaction between two electron spins was measured directly for the first time only recently [21], at a previously inaccessible length scale of $r = 2.4 \mu\text{m}$. Here we exploit this recent measurement to constrain exotic spin-dependent interactions between

electrons, considerably improving on previous bounds at the micrometer scale.

The new measurement reported in Ref. [21] benefits from the high controllability of trapped ions and the resulting relatively straightforward analysis of systematic errors. Briefly, two $^{88}\text{Sr}^+$ ions are trapped in a harmonic potential [$\omega_{\text{trap}} = 2\pi \times 2.386(2)$ MHz] using a linear radio-frequency Paul trap. After laser cooling, the ions form a Coulomb crystal with a separation of $r = 2.407(1) \mu\text{m}$. Each $^{88}\text{Sr}^+$ ion has a single valence electron. A magnetic field of ~ 0.47 mT sets the quantization axis along the line connecting the two ions. The two electrons' state is initialized to $|\uparrow\downarrow\rangle$ with $P_{\text{init}} > 0.98$ fidelity, where $\uparrow(\downarrow)$ indicates a spin polarized along the positive(negative) magnetic field direction. The spins evolve under the spin-spin interaction for $T = 15$ s, ideally resulting in an entangled state $|\Psi(T)\rangle = \cos(2\xi T)|\uparrow\downarrow\rangle + i \sin(2\xi T)|\downarrow\uparrow\rangle$. Finally, the coherence between $|\uparrow\downarrow\rangle$ and $|\downarrow\uparrow\rangle$ is quantified by rotating the spins collectively $|\uparrow\rangle \mapsto (|\uparrow\rangle + |\downarrow\rangle)/\sqrt{2}$, $|\downarrow\rangle \mapsto (|\uparrow\rangle - |\downarrow\rangle)/\sqrt{2}$ and measuring the parity observable, $\Pi \equiv P_{\uparrow\uparrow} + P_{\downarrow\downarrow} - P_{\uparrow\downarrow} - P_{\downarrow\uparrow}$, where P_{ij} represents the probabilities to measure the spin system in respective states, $|ij\rangle$. Since the parity observable is linear in $\sin(4\xi T)$, the coupling strength ξ can be extracted: $\xi = 2\pi \times 1.020(95)_{\text{stat}}$ mHz, where the error is dominated by statistical uncertainty. This approach allows the experimental observable to be insensitive to spatially homogeneous magnetic field noise. The experiment also employed a spin-echo pulse technique [22] to reduce the effect of magnetic field gradients to negligible levels. The measurement of ξ thus gives the interaction strength between the two bound valence electrons of two separate $^{88}\text{Sr}^+$ ions.

The gyromagnetic ratio g of these bound electrons, in the presence of the field of the nucleus and the core electrons, differs from that of free electrons, g_{free} , by $(g_{\text{free}} - g)/g_{\text{free}} \sim 1.8 \times 10^{-5}$ (Breit formula [23]). Therefore, the fractional difference between the measured result and the result that would be obtained for two free electrons can be estimated to be 3.55×10^{-5} , which is well below the measurement statistical uncertainty. This is also the case for all other considered systematic errors in the experiment, which are analyzed and detailed in the supplementary information of Ref. [21].

The measured magnetic interaction strength can easily be compared with the calculated magnetic dipole-dipole interaction between two electrons,

$$\xi_{\text{theory}} = \frac{\mu_0}{4\pi\hbar} \frac{\mu_B^2}{r^3} \left(\frac{g}{2}\right)^2, \quad (1)$$

where μ_0 is the magnetic permeability of vacuum and μ_B is the Bohr magneton. This leads to a theoretical estimate of $\xi_{\text{theory}} = 2\pi \times 0.931(1)$ mHz. The main source of error in determining ξ_{theory} is the statistical uncertainty in r which is estimated from the measured trap frequency, ω_{trap} . Thermal fluctuations of r average out to leading order, since they occur at the trap frequency, and are therefore negligible. The agreement of the measured value $\xi = 2\pi \times 1.020(95)_{\text{stat}}$ mHz with the predicted value ξ_{theory} stands at $\Delta\xi/2\pi = 208$ μ Hz at the 90% confidence level.

This small uncertainty in radial frequency, $\Delta\xi$, translates into new constraints for the strength of axial-vector-mediated interactions between electrons. According to Dobrescu and Mocioiu [1], the exchange of a new vector or axial vector A between fermions results in a Yukawa-type potential [1,2],

$$\mathcal{V}_A(r) = \frac{g_A^e g_A^e \hbar c}{4\pi\hbar c} \frac{1}{r} \mathbf{S}_1 \cdot \mathbf{S}_2 e^{-r/\lambda}, \quad (2)$$

where $g_A^e g_A^e / (4\pi\hbar c)$ is the dimensionless axial-vector coupling constant between the electrons, $\mathbf{S}_{1,2}$ are the electron spins, and r is the interparticle separation.

The \mathcal{V}_A potential would contribute to the $|\uparrow\downarrow\rangle \leftrightarrow |\downarrow\uparrow\rangle$ coherent oscillation measured in Ref. [21], according to the corresponding Hamiltonian, written in the $|\uparrow\downarrow\rangle, |\downarrow\uparrow\rangle$ basis,

$$H_A(r) = \frac{g_A^e g_A^e 2\hbar c}{4\pi\hbar c} \frac{1}{r} e^{-r/\lambda} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (3)$$

The corresponding oscillation frequency,

$$\xi_A = \frac{g_A^e g_A^e c}{4\pi\hbar c} \frac{1}{r} e^{-r/\lambda}, \quad (4)$$

is smaller than $\Delta\xi = 2\pi \times 208$ μ Hz for $r = 2.4$ μ m, at the 90% confidence level. This leads to the constraints plotted in Fig. 1 (dark blue region).

As seen from Eq. (2), the ability to constrain \mathcal{V}_A greatly benefits from the small experimental scale r at which the experiment is performed. Previous measurements performed at scales much greater than a micrometer render much weaker bounds. Therefore, our bounds (dark blue region, Fig. 1) need only be compared to those derivable from smaller scale measurements. Specifically, we compare our result to the measurement of the hyperfine (HF) structure interval of the ground state in positronium (Ps) [26], which constrains exotic dipole-dipole interactions between electrons and positrons above the angstrom scale (light gray region, Fig. 1).

The strength of the axial-vector interaction at the angstrom scale is 4 orders of magnitude larger than at the micron scale due to the $1/r$ dependence of the interaction [Eq. (2)]. Our $^{88}\text{Sr}^+$ based constraints still improve on the Ps measurements due to the exquisite sensitivity of the $^{88}\text{Sr}^+$ experiment, measuring energy shifts with an uncertainty some 10 orders of magnitude smaller than that obtained with Ps spectroscopy. To see this, we note that the present agreement between experimental measurements [24,27,28], $\Delta E_{\text{HF}}(\text{Ps})_{\text{expt}} = 203394.2(1.6)_{\text{stat}}(1.3)_{\text{sys}}$ MHz, and the most recent theoretical calculations based on quantum electrodynamics (QED) [25], $\Delta E_{\text{HF}}(\text{Ps})_{\text{theory}} = 203391.90(25)$ MHz, is 5 MHz at the 90% confidence level [29], dominated by the experimental uncertainty.

Similar to Refs. [26,30], we can estimate the shift of positronium's hyperfine structure interval due to $\mathcal{V}_A(r)$ using first-order perturbation theory. In this case, one must average over the spherically symmetric ground state (see,




FIG. 1 (color online). Constraints (at the 90% confidence level) on the dimensionless coupling constants, $g_A^e g_A^e / (4\pi\hbar c)$ and $g_A^e \bar{g}_A^e / (4\pi\hbar c)$, as a function of the corresponding boson mass. Filled areas correspond to excluded values. The solid line and dark blue fill show the constraints derived from the possible contribution of an axial-vector-mediated interaction to the dipole-dipole interaction between electrons measured in Ref. [21]. The dashed line and light gray region show the constraints on the strength of axial-vector-mediated electron-positron interaction, $g_A^e \bar{g}_A^e / (4\pi\hbar c)$, by comparing the measurement of the ground-state hyperfine interval for positronium [24] to recent QED calculations [25].

for example, Refs. [31–33]) $|\psi\rangle = e^{-r/2a_0}/\sqrt{8\pi a_0^3}$, where a_0 is the Bohr radius. We obtain for the associated energy shift of the Ps ground-state hyperfine structure interval,

$$\Delta E_A = \frac{g_A^e \bar{g}_A^e \hbar c}{4\pi \hbar c 2a_0} \frac{1}{(1 + a_0/\lambda)^2}. \quad (5)$$

As shown in Ref. [30], demanding that $|\Delta E_A| < 2\pi\hbar \times 5$ MHz leads to the constraints shown in Fig. 1 (light gray region). These turn out to be essentially mass independent, since in our region of interest, $\lambda \gg a_0$.

A similar comparison can be made for a pseudoscalar-mediated interaction. A new axionlike pseudoscalar (P) particle results in a dipole-dipole potential between electrons [1,2],

$$\mathcal{V}_P(\mathbf{r}) = \frac{g_P^e g_P^e \hbar^3}{4\pi \hbar c 4m_e^2} \left[\mathbf{S}_1 \cdot \mathbf{S}_2 \left(\frac{1}{\lambda r^2} + \frac{1}{r^3} + \frac{4\pi}{3} \delta^3(r) \right) - (\mathbf{S}_1 \cdot \hat{\mathbf{r}})(\mathbf{S}_2 \cdot \hat{\mathbf{r}}) \left(\frac{1}{\lambda^2 r} + \frac{3}{\lambda r^2} + \frac{3}{r^3} \right) \right] e^{-r/\lambda}, \quad (6)$$

where $g_P^e g_P^e/(4\pi \hbar c)$ is the dimensionless pseudoscalar coupling constant between the electrons, m_e the electron mass, and $\hat{\mathbf{r}}$ is the unit vector along the line connecting the two fermions [34].

Here, the $^{88}\text{Sr}^+$ experimental sensitivity is not sufficient to overcome the larger pseudoscalar interaction strength at the angstrom scale probed by the Ps measurement, due to the $1/r^3$ dependence of the potential. For the $^{88}\text{Sr}^+$ experiment, \mathcal{V}_P would contribute to the $|\uparrow\downarrow\rangle\leftrightarrow|\downarrow\uparrow\rangle$ coherent oscillation measured in Ref. [21] by

$$\xi_P = \frac{g_P^e g_P^e \hbar^2}{4\pi \hbar c 4m_e^2} \left(\frac{1}{r^3} + \frac{1}{\lambda r^2} \right) e^{-r/\lambda}. \quad (7)$$

Demanding that ξ_P be smaller than $\Delta\xi = 2\pi \times 208$ μHz for $r = 2.4$ μm at the 90% confidence level leads to the constraints plotted in Fig. 2 (dark-blue region). In comparison, the energy shift of the Ps ground-state hyperfine structure interval due to $\mathcal{V}_P(r)$ is

$$\Delta E_P = \frac{g_P^e \bar{g}_P^e \hbar^3}{4\pi \hbar c 24m_e^2 c a_0^3} \left(1 - \frac{1}{(1 + \lambda/a_0)^2} \right). \quad (8)$$

As shown in Ref. [26], demanding that $|\Delta E_P| < 2\pi\hbar \times 5$ MHz leads to the constraints depicted by the light gray region in Fig. 2, again, as in Fig. 1, showing nearly mass-independent behavior over the depicted range.

We note that direct comparison between the electron-electron and electron-positron constraints is based on the implicit assumption of *CPT* invariance for the exotic interactions studied here. The potentials $\mathcal{V}_A(r)$ and $\mathcal{V}_P(r)$ are even under the parity (P) and time-reversal (T) transformations, and thus they are also even under the




FIG. 2 (color online). Same as Fig. 1, for the case of a pseudoscalar-mediated interaction, constraining $g_P^e g_P^e/(4\pi \hbar c)$ for electron-electron (dark blue region) and $g_P^e \bar{g}_P^e/(4\pi \hbar c)$ for electron-positron (light gray region) interactions.

combined *PT* symmetry. Consequently, assuming *CPT* invariance (where C represents charge conjugation), electrons and positrons should have the same magnitude of coupling strength to exotic pseudoscalar or axial-vector interactions: $|g_P^e| = |\bar{g}_P^e|$ and $|g_A^e| = |\bar{g}_A^e|$. It should be noted, however, that *CPT* invariance is not guaranteed for these exotic interactions, at least based on the current state of knowledge [35]. Thus, one can regard the e^- - e^- and the e^+ - e^- measurements discussed here as independent constraints.

The above analysis can also be used to place bounds on torsion gravity. According to general relativity, the local space-time curvature is unaffected by the presence of spin [36–40]. However, in extensions of general relativity based on a Riemann-Cartan space-time, the gravitational interaction is described by a torsion tensor which can generate spin-mass and spin-spin interactions [41–46]. The spin-spin interaction generated by a standard propagating torsion field takes the form of $\mathcal{V}_P(r)$ [47,48], with $\lambda \rightarrow \infty$ and a coupling strength that can be parametrized in terms of a dimensionless parameter β , such that

$$\beta^2 = \left(\frac{g_P^e g_P^e}{4\pi \hbar c} \right) \left(\frac{2 \hbar c}{9 G m_e^2} \right), \quad (9)$$

where G is the gravitational constant. The minimally coupled Dirac equation for a spin-1/2 particle [47,49–51] predicts $\beta = 1$. Experimental constraints, based on the analysis presented here, are shown in Table I.

Finally, we note that stellar energy-loss arguments strongly constrain the pseudoscalar coupling of electrons [56], in particular, revealing that for particles with mass $m \lesssim 10$ keV [57], $|g_P^e g_P^e/(4\pi \hbar c)| \lesssim 10^{-25}$, far exceeding the laboratory limits discussed here. In terms of torsion gravity, these astrophysical constraints translate to a limit $\beta^2 \lesssim 10^{19}$. These constraints, however, do not apply to the axial-vector interactions [1].

TABLE I. Comparison of torsion gravity constraints.

| Particles | Experiment | Scale \geq | $ \beta^2 \leq$ | Reference |
|------------|------------|-----------------|--------------------|-----------------------------|
| $n-n$ | | 50 cm | 2×10^{28} | [52] |
| $p-p$ | | 50 cm | 2×10^{31} | [53] (experiment [52]) |
| e^-e^- | | 15 cm | 4×10^{28} | [19] |
| e^-e^- | | 2 μm | 2×10^{41} | this work (experiment [21]) |
| $n-n$ | | 100 nm | 6×10^{34} | [54] |
| $p-p$ | | \AA | 1×10^{33} | [55] |
| e^-e^+ | | \AA | 3×10^{38} | this work (experiment [24]) |
| $p-p, n-p$ | | \AA | 1×10^{31} | [32] |

In conclusion, the results of a new direct measurement of the magnetic interaction between two electrons at the micrometer scale were used to place bounds on exotic forces. The constraints on an axial-vector-mediated spin-spin interaction between electrons are 6 orders of magnitude more stringent than those derived from positronium spectroscopy for electron-positron pairs, at the micrometer scale (equivalently for masses below ~ 1 eV). Note that the constraint could be improved by 1 or 2 orders of magnitude, based on an extension of the technology used in Ref. [21]. This would require a tighter ion trap, consequently placing the two electronic spins at separations less than a micrometer.

The collaboration resulting in this paper was suggested by Dmitry Budker. The authors are sincerely grateful to Mitch Watnik, Joshua Kerr, Gilad Perez, Moty Milgrom, and Guy Gur-Ari Krakover, for enlightening discussions. We thank Dmitry Budker and Jennie Guzman for their careful reading of our manuscript. This work was supported, in part, by the U.S. National Science Foundation under Grant No. PHY-1307507 and by I-Core: The Israeli Excellence Center “Circle of Light” and the European Research Council (consolidator Grant No. Ionology-616919).

- [1] B. A. Dobrescu and I. Mocioiu, *J. High Energy Phys.* **11** (2006) 005.
- [2] J. E. Moody and F. Wilczek, *Phys. Rev. D* **30**, 130 (1984).
- [3] S. Weinberg, *Phys. Rev. Lett.* **40**, 223 (1978).
- [4] F. Wilczek, *Phys. Rev. Lett.* **40**, 279 (1978).
- [5] R. D. Peccei and H. R. Quinn, *Phys. Rev. Lett.* **38**, 1440 (1977).
- [6] B. A. Dobrescu, *Phys. Rev. Lett.* **94**, 151802 (2005).
- [7] T. Appelquist, B. A. Dobrescu, and A. R. Hopper, *Phys. Rev. D* **68**, 035012 (2003).
- [8] G. Bertone, D. Hooper, and J. Silk, *Phys. Rep.* **405**, 279 (2005).
- [9] A. Friedland, H. Murayama, and M. Perelstein, *Phys. Rev. D* **67**, 043519 (2003).
- [10] V. Flambaum, S. Lambert, and M. Pospelov, *Phys. Rev. D* **80**, 105021 (2009).

- [11] P. W. Graham, D. E. Kaplan, and S. Rajendran, [arXiv:1504.07551](https://arxiv.org/abs/1504.07551).
- [12] A. Arvanitaki, S. Dimopoulos, S. Dubovsky, N. Kaloper, and J. March-Russell, *Phys. Rev. D* **81**, 123530 (2010).
- [13] P. V. Vorobyov and Ya. I. Gitarts, *Phys. Lett. B* **208**, 146 (1988).
- [14] R. C. Ritter, C. E. Goldblum, W.-T. Ni, G. T. Gillies, and C. C. Speake, *Phys. Rev. D* **42**, 977 (1990).
- [15] V. F. Bobrakov, Yu. V. Borisov, M. S. Lasakov, A. P. Serebrov, R. R. Tal'daev, and A. S. Trofimova, *Pis'ma Zh. Eksp. Teor. Fiz.* **53**, 283 (1991) [JETP Lett. **53**, 294 (1991)].
- [16] D. J. Wineland, J. J. Bollinger, D. J. Heinzen, W. M. Itano, and M. G. Raizen, *Phys. Rev. Lett.* **67**, 1735 (1991).
- [17] T. C. P. Chui and W. T. Ni, *Phys. Rev. Lett.* **71**, 3247 (1993).
- [18] L. Hunter, J. Gordon, S. Peck, D. Ang, and J.-F. Lin, *Science* **339**, 928 (2013).
- [19] B. R. Heckel, W. A. Terrano, and E. G. Adelberger, *Phys. Rev. Lett.* **111**, 151802 (2013).
- [20] As can be seen from Eq. (2) and Eq. (6), extrapolating these bounds to the micrometer scale deteriorates their sensitivity roughly by a factor of e^x/x , where x is the ratio between the measurement scale and the extrapolated scale, i.e., $x \geq 1 \text{ mm}/1 \mu\text{m} = 1000$. That renders bounds obtained at the millimeter scale useless at the micrometer scale.
- [21] S. Kotler, N. Akerman, N. Navon, Y. Glickman, and R. Ozeri, *Nature (London)* **510**, 376 (2014).
- [22] S. Kotler, N. Akerman, Y. Glickman, A. Keselman, and R. Ozeri, *Nature (London)* **473**, 61 (2011).
- [23] G. Breit, *Nature (London)* **122**, 649 (1928).
- [24] A. Ishida, T. Namba, S. Asai, T. Kobayashi, H. Saito, M. Yoshida, K. Tanaka, and A. Yamamoto, *Phys. Lett. B* **734**, 338 (2014).
- [25] M. I. Eides and V. A. Shelyuto, *Phys. Rev. D* **89**, 111301(R) (2014).
- [26] T. M. Leslie, E. Weisman, R. Khatiwada, and J. C. Long, *Phys. Rev. D* **89**, 114022 (2014).
- [27] A. P. Mills, Jr. and G. H. Beaman, *Phys. Rev. Lett.* **34**, 246 (1975); A. P. Mills, Jr., *Phys. Rev. A* **27**, 262 (1983).
- [28] M. W. Ritter, P. O. Egan, V. W. Hughes, and K. A. Woodle, *Phys. Rev. A* **30**, 1331 (1984).
- [29] Statistical and systematic uncertainties have been combined in quadrature and a normal distribution has been assumed.
- [30] S. G. Karshenboim, *Phys. Rev. Lett.* **104**, 220406 (2010); *Phys. Rev. D* **82**, 073003 (2010); *Phys. Rev. D* **82**, 113013 (2010).
- [31] D. F. Jackson Kimball, A. Boyd, and D. Budker, *Phys. Rev. A* **82**, 062714 (2010).
- [32] M. P. Ledbetter, M. V. Romalis, and D. F. Jackson Kimball, *Phys. Rev. Lett.* **110**, 040402 (2013).
- [33] W.-Y. Keung, and D. Marfatia, [arXiv:1501.00455](https://arxiv.org/abs/1501.00455).
- [34] Note that Ref. [1] neglects the term containing the delta function, since Dobrescu and Mocioiu focus on long-range potentials; the delta function term is present in the analogous equation from Ref. [2].
- [35] B. Altschul, *Phys. Rev. D* **87**, 045012 (2013).
- [36] I. Y. Kobzarev and L. B. Okun, *Zh. Eksp. Teor. Fiz.* **43**, 1904 (1962) [Sov. Phys. JETP **16**, 1343 (1963)].
- [37] J. Leitner and S. Okubo, *Phys. Rev.* **136**, B1542 (1964).

[38] F. W. Hehl and W.-T. Ni, *Phys. Rev. D* **42**, 2045 (1990).

[39] I. B. Khriplovich and A. A. Pomeransky, *J. Exp. Theor. Phys.* **86**, 839 (1998) [Zh. Eksp. Teor. Fiz. **113**, 1537 (1998)].

[40] A. J. Silenko and O. V. Teryaev, *Phys. Rev. D* **71**, 064016 (2005).

[41] F. W. Hehl, P. von der Heyde, G. D. Kerlick, and J. M. Nester, *Rev. Mod. Phys.* **48**, 393 (1976).

[42] I. L. Shapiro, *Phys. Rep.* **357**, 113 (2002).

[43] R. T. Hammond, *Rep. Prog. Phys.* **65**, 599 (2002).

[44] V. A. Kostelecky, N. Russell, and J. D. Tasson, *Phys. Rev. Lett.* **100**, 111102 (2008).

[45] Y. N. Obukhov, A. J. Silenko, and O. V. Teryaev, *Phys. Rev. D* **90**, 124068 (2014).

[46] R. Lehnert, W. M. Snow, and H. Yan, *Phys. Lett. B* **730**, 353 (2014).

[47] R. T. Hammond, *Phys. Rev. D* **52**, 6918 (1995).

[48] E. G. Adelberger, J. H. Gundlach, B. R. Heckel, S. Hoedl, and S. Schlammung, *Prog. Part. Nucl. Phys.* **62**, 102 (2009).

[49] D. E. Neville, *Phys. Rev. D* **21**, 2075 (1980).

[50] D. E. Neville, *Phys. Rev. D* **25**, 573 (1982).

[51] S. M. Carroll and G. B. Field, *Phys. Rev. D* **50**, 3867 (1994).

[52] G. Vasilakis, J. M. Brown, T. W. Kornack, and M. V. Romalis, *Phys. Rev. Lett.* **103**, 261801 (2009).

[53] D. F. Jackson Kimball, arXiv:1407.2671.

[54] C. Fu and W. M. Snow, arXiv:1103.0659.

[55] N. F. Ramsey, *Physica (Amsterdam)* **96A**, 285 (1979).

[56] G. Raffelt, *Phys. Rev. D* **86**, 015001 (2012).

[57] G. Raffelt and A. Weiss, *Phys. Rev. D* **51**, 1495 (1995).