

Spatial Relationships of Microtubule-organizing Centers and the Contact Area of Cytotoxic T Lymphocytes and Target Cells

BENJAMIN GEIGER,* DALIA ROSEN, and GIDEON BERKE

Departments of Chemical Immunology and Cell Biology, The Weizmann Institute of Science, Rehovot 76100, Israel*

ABSTRACT Specific binding (conjugation) of cytotoxic T lymphocytes (CTL) to target cells (TC) is the first step in a multistage process ultimately resulting in dissolution of the TC and recycling of the CTL. We examined the position of the microtubule organizing center (MTOC) of immune CTL bound to specific TC. Immunofluorescence labeling of freshly prepared CTL-TC conjugates with tubulin antibodies indicated that the MTOC in essentially all conjugated CTL but not in the conjugated TC were oriented towards the intercellular contact site. This finding was corroborated by electron microscopy examination of CTL-TC conjugates fixed either immediately after conjugation or during the lytic process. Antibody-induced caps of membrane antigens of CTL such as H-2 and Thy 1, did not show a similar relationship to the MTOC. Incubation of CTL-TC conjugates, 10–15 min at room temperature, resulted in an apparent deterioration of the microtubular system of conjugated CTL. It is proposed that the CTL plasma membrane proximal to the MTOC is particularly active in forming stable intercellular contacts, resulting in CTL-TC conjugation, and that subsequent modulation of the microtubular system of the CTL may be related to the cytolytic response and to detachment of the effector cell.

A prominent manifestation of cell-mediated immunity is the lytic interaction of cytotoxic T lymphocytes (CTL) with appropriate target cells (TC). This process is believed to be relevant to virus, tumor, and transplantation immunity (4, 13, 42). The first step in CTL-mediated lysis is the binding of CTL and TC (conjugation) mediated through specific CTL cell surface receptor(s) and TC major histocompatibility complex determinants. Binding is followed by a lethal hit step delivered by the CTL, ultimately leading to lysis. Following lysis of the TC, effector CTL detach and can recycle to start a new lytic interaction (see references 4, 7, 8, 18, 19, 23, 28 for reviews).

Several observations suggest that the CTL are polar from at least a functional point of view. It has been shown that the lethal hit is strictly unidirectional, i.e., it affects only CTL-conjugated TC without causing damage to the effector CTL (40). This "immunity" of the CTL cannot be attributed to an inherent resistance towards the cytolytic process, since CTL of a given type can be readily killed by other specifically immunized CTL (14, 17). Moreover, it has been observed that although an individual CTL can bind a number of TC simultaneously, lysis of individual TC occurs sequentially (40, 41). These and other (17) results suggest that the lethal hit, whatever its nature, is expressed in a polar, unidirectional fashion.

In this study we examine the possibility that the unidirectional killing activity of CTL is related, at least in part, to a specific polar arrangement of the cytoskeletal system of either the CTL or the TC. We present evidence suggesting polarity by showing that CTL bind predominantly through a cell surface region proximal to the microtubule organizing center (MTOC). Moreover, we demonstrate that following CTL-TC binding, the CTL microtubular system becomes partially deteriorated. The significance of these findings to the mechanism of cytotoxic interactions is discussed.

MATERIALS AND METHODS

Animals and Tumors

Highly inbred, C57BL/6 (major histocompatibility H-2^b) and BALB/c (H-2^d) mice were provided by the Weizmann Institute animal-breeding center. Leukemia EL4 and mastocytoma P815 cells were maintained in ascitic form by weekly transfer of about 25×10^6 cells into syngeneic recipients (C57BL/6 and DBA/2 respectively).

Generation of CTL

A system to generate and study alloimmune peritoneal exudate CTL has been described before (5). Briefly, 10–12 d after intraperitoneal injection of 25×10^6 EL4 leukemia cells, BALB/c anti-EL4 mice were killed by CO₂ narcosis and

their peritoneal cavities rinsed with PBS supplemented with 10% fetal calf serum (PBS-FCS). The peritoneal cells were centrifuged 10 min at 250 g at 4°C, resuspended in 15 ml PBS-FCS per four to seven mice, and incubated on nylon wool columns at 37°C for 45 min (5) to remove adherent cells (macrophages and others). Nonadherent cells containing 30–40% (4) CTL were eluted by rinsing the columns with 20 ml cold PBS-FCS.

CTL-TC Binding (Conjugation)

The method involved centrifugation of mixtures containing CTL and TC, leading to formation of CTL-TC clusters, referred to as conjugates, which can be examined and studied individually under the microscope (6). 1 ml suspensions containing 10^6 CTL and 10^6 TC in PBS-FCS, in 12 × 75 mm tubes were centrifuged at 170 g for 10 min at room temperature. The pellets were resuspended vigorously with a Pasteur pipette and the number of CTL-TC conjugates was counted in a hemocytometer.

CTL-mediated Lysis of TC

The chromium-release technique (12, 13) was used to measure the cytolytic capacity of CTL. EL4 or P815 TC were removed from the peritoneal cavity of syngeneic hosts and 3×10^7 cells in 1 ml PBS-FCS were incubated with 200 μ Ci $\text{Na}_2^{51}\text{CrO}_4$ (Amersham CJS, I.P., Amersham Corp., Arlington Heights, IL) for 45 min at 37°C with occasional shaking, and washed twice with PBS-FCS. Labeled TC (1×10^5 cells) were mixed with CTL (3×10^6 cells) in 1-ml aliquots in 75 × 12 mm test tubes, centrifuged at 170 g for 10 min to promote CTL-TC contact, and incubated at 37°C in a humidified atmosphere. At the end of the incubation period the radioactivity of the supernatant was assayed in a well-type gamma scintillation counter. Released radioactivity is expressed as the percent of total releasable radioactivity, determined by repeated freezing and thawing of labeled target cells. Results are corrected for spontaneous release (10–15% of total radioactivity).

Electron Microscopy

BALB/c anti-EL4 CTL (1×10^6) were mixed with EL4 TC ($1-3 \times 10^5$) in 1 ml PBS-FCS in Falcon (Oxnard, CA) 75 × 12-mm plastic tubes at room temperature. Cells were centrifuged at 170 g for 10 min at room temperature to promote CTL-TC conjugate formation. Medium was removed and carefully replaced by prewarmed (37°C) fixative (2.5% glutaraldehyde in 0.09 M cacodylate buffer, pH 7.2, containing 2.5 mM CaCl_2). Prewarming of the fixative was important to prevent retraction of cellular projections due to temperature change and for the preservation of intact microtubules. After 15 min at 37°C, and an additional 45 min at room temperature, the cells were washed overnight with the same buffer, postfixed for 1 h in 1% osmium tetroxide, washed briefly with distilled water, stained in block with 0.5% aqueous uranyl acetate for 1 h, dehydrated in ethanol and embedded in Poly/Bed 812 (Polysciences, Inc., Paul Valley Industrial Park, Warrington, PA). Thin sections were cut on Sorvall MT-2 ultramicrotome (New Town, CT), stained with uranyl acetate and lead citrate, and examined in a Philips EM300 electron microscope operated at 80 kV. Conjugates were examined preferentially at the binding site area.

Immunofluorescence

Centrifuged pellets of CTL/TC mixtures containing 1×10^6 of each cell type were suspended in 1 ml PBS and drops were applied to polylysine-coated glass cover slips. After 2–3 min at room temperature, cover slip-adherent cells were fixed in 3% paraformaldehyde for 30 min. The attached cells including single cells as well as conjugates were permeabilized with 0.2% Triton X-100 (in PBS) for 4 min and indirectly labeled for tubulin using affinity-purified rabbit antitubulin antibodies followed by rhodamine-conjugated goat anti-rabbit IgG, as previously described (16). Fluorescence microscopy was performed using a Zeiss photomicroscope III equipped with epilluminator and filter sets for fluorescein and rhodamine. Photomicrographs were usually taken through an ×63/1.4 planapochromat objective (suitable for fluorescence and Nomarski optics) or through an ×63/1.25 Neofluar objective (suitable for phase-contrast optics as well).

Capping Experiments

Antiserum specific to H-2^d (C57BL/6 anti-P 815) was diluted 1:20 in PBS and incubated for 15 min at room temperature with BALB/c anti-EL4 CTL. After rinsing, the cells were mixed with fluorescein-conjugated goat anti-mouse IgG (affinity purified, 20 μ g/ml) and incubated for 20 min at 37°C. The cells were then washed, plated on polylysine-coated cover slips and fixed. Capping of Thy-

1 antigen was carried out in a similar manner, using a 1:100 dilution of monoclonal antibody solution (ascitic fluid) reactive with this antigen (22).

RESULTS

Lysis of EL4-TC by Peritoneal BALB/c Anti-EL4 CTL

To ascertain that the CTL studied below are active effector cells capable of binding to and killing TC, the kinetics of their binding to and lysis of TC were studied (Fig. 1).

After centrifugation of CTL/TC mixtures containing 1×10^6 BALB/c anti-EL4 cells (CTL) and 3.3×10^5 EL4 (TC), ~46% of the TC were found conjugated to CTL. Upon incubation at 37°C, lysis of the TC occurred, and the percentage of lysed TC, as monitored by the release of chromium, increased with time resulting in a progressive decrease in the amount of conjugates. (After 60 min of incubation only 14.5% conjugated TC could be detected).

The Spatial Relationships of MTOC and CTL-TC Contact Areas

IMMUNOFLUORESCENCE MICROSCOPY: Specific conjugates of BALB/c anti-EL4 CTL and EL4 cells (TC) were prepared by centrifuging mixtures containing CTL and TC at room temperature, as described (6). Microscopic examination of the cells with differential interference contrast or phase-contrast optics revealed a large number of CTL-TC conjugates comprised of one CTL bound to one TC (Fig. 2 C). EL4 cells (average diameter of 15–20 μ m) could easily be distinguished from the much smaller CTL (7–10 μ m) (5, 6). The microtubular system of nonconjugated (free) CTL consisted of about 10–30 visible, fluorescently-labeled microtubules or microtubule bundles. Careful examination of the labeled cells revealed a single microtubule organizing center (MTOC) in essentially every cell (Fig. 2 E). EL4 cells exhibited a dense microtubular network (Fig. 2 F) originating in a single MTOC. In CTL-TC conjugates we found that essentially all effector CTL were bound to EL4 through a membrane region proximal to the MTOC. This polar orientation was observed in over 95% of the cells fixed immediately after onset of conjugation by centrifugation (out of over 500 conjugated cells analyzed). Conjugated TC on the other hand, exhibited a random orientation of their contact zone with respect to the MTOC. A single CTL-TC conjugate is




FIGURE 1 Conjugation and cytolysis of EL4 TC by BALB/c anti-EL4 CTL. Cells (CTL 3:TC 1) were mixed, centrifuged 170 g for 10 min at room temperature, and incubated at 37°C. At various times thereafter percent lysis was monitored by the ^{51}Cr -release technique and percent conjugation was determined microscopically. Data based on five independent experiments. Solid line, % ^{51}Cr released. Broken line, % conjugation.










FIGURE 2 Indirect immunofluorescence labeling for tubulin of CTL-TC conjugates. The cells were fixed immediately after centrifugation. (A and B) Tubulin immunolabeling of the same conjugate, photographed at two focal planes to show the MTOC of the CTL (smaller cell) and the TC (larger cell), respectively. (C) The same conjugate, photographed with Nomarski optics. (D) Immunofluorescent labeling for tubulin of conjugates consisting of one TC and two or three CTL. Notice that in all cases the MTOC of the CTL is proximal to the area of contact, while the MTOC of the TC apparently has a random orientation. (E) Unconjugated CTL. (F) Unconjugated EL4 TC. Bars, 10 μ m. A-C, E, F: \times 900. D, \times 1,200.

shown in Fig. 2A and 2B. The cells were photographed at two focal planes showing the MTOC of the effector CTL (Fig. 2A) and of the TC (Fig. 2B).

At effector cell excess we frequently observed complex conjugates consisting of a number of effector cells bound to one target (see also reference 6). In Fig. 2D, three types of conjugates (CTL 1-TC 1; CTL 2-TC 1; CTL 3-TC 1) are shown. In these, the MTOC of all bound CTL were oriented towards the target, while the MTOC of the latter was randomly distributed.

ELECTRON MICROSCOPY: The results outlined above were corroborated by transmission electron microscopy. Examination of a large number of CTL-TC conjugates, sectioned

across intercellular contact sites, indicated that the centrioles as well as the pericentriolar array of microtubules of the CTL were located proximal to the contact area. This was found in conjugates at early stages of the cytotoxic interaction (Fig. 3A, B, and F) or in relatively late stages where the damage to the TC was clearly apparent (Fig. 3C and D). Due to the extensive membrane interdigitation at the contact area and the random plane of sectioning (with respect to the contact plane) we could not determine whether the axes of the centrioles themselves were oriented toward the cell membrane. In high magnifications (Fig. 3B, D, and F) we could detect a large number of microtubules emerging from the centriole complex. In some






FIGURE 4 Double immunofluorescence-labeling of CTL for surface patches and caps induced by anti-H-2^d (A) or anti-Thy-1 (D) and for tubulin (B and E, respectively). The same microscopic fields photographed with Normarski optics are shown in C and F. Arrowheads point to the polar caps induced by the antibodies, and the arrows to the MTOC of the same cells. Bar, 10 μ m. \times 600.

CTL we detected elements of the Golgi complex in the vicinity of the contact area in the CTL (for example the cell shown at the top of Fig. 3E and in Fig. 3F). Nevertheless, this was not nearly as prominent as the presence of the MTOC in that region.

The centrioles of the EL4 TC were apparently randomly distributed with respect to the intercellular contact area as demonstrated in Fig. 3E. It should be pointed out that the number of MTOC detected in CTL in ultrathin sections selected for the CTL-TC contact area was considerably higher (more than twofold) than that detected in unselected sections or unconjugated cells.

Ligand-induced Surface Caps are not Spatially Related to the MTOC

Incubation of BALB/c anti-EL4 CTL carrying both the H-2^d and Thy-1 antigens with H-2^d and Thy-1 antibodies followed by rhodamine-labeled goat anti-mouse IgG resulted in redistribution of the respective molecules into patches and subsequently into polar caps (Fig. 4A and D). Usually 20–30 min of incubation at 37°C were required for maximal cap formation.

Capped, fixed cells were permeabilized with Triton X-100 and reacted with a rabbit antitubulin antibody followed by fluorescein-labeled goat anti-rabbit IgG. The results (Fig. 4) indicate that the polar caps formed in the two systems (A and D) were not spatially related to the MTOC (B and E). Moreover, in most cases (ca. 60–75%) the MTOC were localized opposite to the cap. Comparison of the microtubular system of cells before and after capping suggested that some deterioration of the microtubule network occurred during capping (compare for example Fig. 4B and E with Fig. 2E).

Modulation of the Microtubular System in TC-bound CTL

CTL-mediated lysis is highly temperature-dependent; incubation of CTL-TC conjugates at room temperature resulted in virtually no killing (6). Nevertheless, we noticed that after 5–10 min incubation at room temperature (or at 37°C) the microtubule system of TC-bound CTL underwent substantial deterioration. This was usually manifested by a decrease in number or complete disappearance of defined microtubules (Fig. 5). Documentation of this effect required examination of the cells

FIGURE 3 Electron microscopy of CTL-TC conjugates. (A) Low-power magnification of one target cell (TC) bound to two CTL (top left and bottom). In the upper CTL the cross-sectioned centriole is clearly localized near the contact region (arrow). (B) CTL-TC contact area showing the centriole pair in the CTL (marked with two arrows). The target cell (TC) appears intact. (C and D) Low-power (C) and high-power (D) magnifications of conjugate consisting of two CTL bound to one damaged target cell (TC). D is an enlarged portion of C (the asterisk in both photographs is in identical position). The two CTL are bound to the TC with their MTOC (arrows) proximal to the contact zone. In D, many microtubules can be seen emerging from the pericentriolar region. (E) Electron photomicrograph showing a CTL-TC conjugate. In the CTL, the centrioles or pericentriolar microtubules (arrows) as well as the Golgi apparatus (arrowhead) are oriented towards the contact area, while the centriole of the TC is not. (F) Longitudinal section through the centriole (arrow) with surrounding microtubules. Part of the Golgi system (arrowhead) and vacuoles are observed in the vicinity of the CTL membrane at the contact area. Bars, 0.1 μ m. A, \times 70,000; B, \times 220,000; C, \times 55,000; D, \times 120,000; E, \times 45,000; F, \times 270,000.




FIGURE 5 Indirect immunofluorescence labeling for tubulin of conjugates, fixed after 15 min at room temperature. The conjugate in *A* and *B* was photographed at two focal planes to demonstrate that whereas the MTOC can still be detected near the contact area, the microtubular system of the CTL appears deteriorated. (*C*) Unconjugated CTL which was exposed to the same incubation. Bar, 10 μ m. \times 950.

at several focal planes. No changes were noted in the organization of the microtubular systems in the target cells nor were such changes observed in nonconjugated CTL.

DISCUSSION

Extensive efforts were invested in recent years to characterize the mechanisms involved in the interactions of cytotoxic T lymphocytes with specific targets (for reviews see 4, 13, 18, 19, 23, 28). Many studies focused on the early steps of the process and the nature of the CTL receptor for TC recognition, whereas others explored the lytic process itself. Although the cytotoxic mechanism of CTL has not yet been defined at the molecular level, several models have been considered to account for the lethal hit, ranging from secretion of toxic substances, through mechanical deformation, to induction of local instabilities in the TC membrane. Whichever working hypothesis better presents the cytotoxic effect, it must be compatible with a fundamental feature of this process, that is, the unidirectionality of the killing. It has been established that after the target cell has been damaged, the CTL can recycle to start a new lytic interaction (40). The killer cell, however, is not inherently immune to killing, as it has been shown that CTL can be killed

when exposed to appropriately immunized effector cells (17).

A finding related to unidirectional killing by CTL is that effector CTL bound simultaneously to more than one TC kill the TC sequentially (41). This suggested that the delivery of the lethal hit not only requires direct intercellular contact but preferentially occurs at defined regions along the CTL plasma membrane. Electron microscopy examination of CTL-TC conjugates revealed extensive interdigitation of the plasma membranes at the contact region (26, 29), suggesting that membrane-folding forces are generated at CTL-TC contact zones. In view of the involvement of cytoskeletal elements in the mechanical responses of cells (20, 25, 35), we explored the possibility that specific organization of such intracellular networks might be related to the unidirectionality of the cytotoxic process. It has been recently shown by immunofluorescence microscopy that the CTL contact area is enriched with actin (27), in analogy to the increase in actin labeling under lectin- or antibody-induced surface caps (9, 15, 30, 33). Actin enrichment at CTL-TC contact regions is compatible with the extensive membrane interdigitations observed at this site (26, 29).

The observations reported here were related to the microtubule system and its organizing center. They may be summarized as follows: (*a*) immune CTL interact with and bind to TC predominantly through a membrane region proximal to the MTOC and the centrioles; (*b*) unlike CTL, CTL-bound TC do not display their MTOC proximal to the contact region; (*c*) after incubation at room temperature (or 37°C), the microtubules of TC-bound CTL undergo progressive deterioration; (*d*) antibody-induced caps of H-2 and Thy-1 antigens do not localize proximal to the MTOC; often they are found in the opposite pole of the cell.

The two phenomena, namely polar orientation of the MTOC and deterioration of microtubules seem to be related to distinct, sequential events and will be considered separately. The results indicate that the plasma membrane proximal to the MTOC is engaged in binding to the target, at least in conjugates composed of one CTL bound to one TC. This spatial relationship was found in essentially all CTL-TC conjugates fixed immediately after onset of conjugation. It is still unclear whether it is related to an uneven distribution of relevant membrane components or to adhesive properties of the cell membrane adjacent to the MTOC which favor the formation of stable intercellular contacts. It may be related to the more general role of the centriole in cellular dynamics (for discussion see references 1, 2). It has been shown that in mobile cells the centrioles are localized in front of the nucleus, towards the leading edge of the cell membrane (3, 21). The membrane in this area exhibits an increased protrusive and deformational potential which may render it more compatible for the formation of stable intercellular contacts.

Another organelle found localized in the vicinity of CTL contact regions is the Golgi complex. This is in line both with reports by Zagury et al. (40) and by Bykovskaya et al. (10, 11) and with the notion that the Golgi apparatus is usually localized in the vicinity of the centrioles (32, 34). However, analysis of large numbers of electron micrographs of CTL-TC conjugates suggests that the contact area is primarily related to the centrioles and that the localization of the Golgi system is only secondarily related to the contact site.

Patching or capping of surface components on the CTL plasma membrane which might occur during CTL-TC interaction could not in itself account for the orientation of the MTOC inasmuch as surface caps induced by H-2 or Thy 1, antibodies, and rhodamine-labeled goat anti-mouse IgG were

not proximal to the MTOC and were often localized at the opposite pole. It was shown (31) that capping of these antigens occurred predominantly in the area opposite the Golgi region (in contrast to anti-IgG or Con A induced caps [24, 31, 36]).

The second phenomenon reported above was the deterioration of the microtubule network of conjugated effector cells. This process, though difficult to evaluate quantitatively, was quite rapid and occurred progressively after incubation of cell conjugates for 5–15 min at room temperature or at 37°C. Similar modulation of microtubule organization was often found in cells that undergo patch or cap formation (Fig. 4). These observations are in line with the report of Yahara and Kakimoto-Sameshima (39) on the modulation of microtubule organization by capping of surface Ig in mouse spleen lymphocytes. Unlike CTL-TC conjugation which requires Mg^{++} , the lytic process is strictly Ca^{++} dependent (23). However, it is still unclear whether this Ca^{++} dependence is related to well-known effects of Ca^{++} on microtubule disassembly.

The significance of modulation of microtubule organization in CTL and its relevance to the cytotoxic response are at present not clear. One may consider, however, the distinct and often conflicting actions of microtubules and actin-containing microfilaments in living cells. It has been shown that paralysis of the lateral mobility of surface components induced by high concentrations of Con A could be abolished by the microtubule-disrupting drug colchicine (37, 38). It may be proposed that the intact network of microtubules restrains dynamic, actin-dependent processes of the membrane such as those that might be involved in cytotoxic interaction. Whether deterioration of microtubule organization in CTL is directly related to an abolition of this restraint, to the activation of the actin-containing contractile system, and/or to the potentiation of a lytic process is yet to be determined.

Benjamin Geiger is an incumbent of the Charles Revson Chair in Biology. Gideon Berke is Issac and Elsa Bourla Professor of Cancer Research.

Received for publication 1 April 1982, and in revised form 19 May 1982.

REFERENCES

- Albrecht-Buehler, G. 1979. Navigation of nonneuronal cells. In *The Role of Intercellular Signals: Navigation, Encounter, Outcome*. J. G. Nicholls, editor. Berlin: Dahlem Konferenzen. 75–96.
- Albrecht-Buehler, G. 1981. Does the geometric design of centrioles imply their function? *Cell Motility*. 1:237–245.
- Albrecht-Buehler, G., and A. Bushnell. 1979. The orientation of centrioles in migrating 3T3 cells. *Exp. Cell Res.* 120:111–118.
- Berke, G. 1980. Interaction of cytotoxic T-lymphocytes and target cells. *Prog. Allergy*. 27:69–133.
- Berke, G., K. A. Sullivan, and D. B. Amos. 1972. Rejection of ascites tumor allografts. I. Isolation, characterization and *in vitro* reactivity of peritoneal lymphoid effector cells from BALB/c mice immune to EL4 leukemia. *J. Exp. Med.* 135:1334–1350.
- Berke, G., D. Gabison, and M. Feldman. 1975. The frequency of effector cells in populations containing cytotoxic T-lymphocytes. *Eur. J. Immunol.* 5:813–818.
- Berke, G., V. Hu, E. McVey, and W. R. Clark. 1981. T lymphocyte-mediated cytotoxicity. I. A common mechanism for target recognition in specific and lectin-dependent cytotoxicity. *J. Immunol.* 127:776–781.
- Berke, G., E. McVey, V. Hu, and W. R. Clark. 1981. T lymphocyte-mediated cytotoxicity. II. Role of target cell histocompatibility antigens in recognition and lysis. *J. Immunol.* 127:782–787.
- Bourguignon, L. Y. W., and S. J. Singer. 1977. Transmembrane interaction and the mechanism of capping of surface receptor by their specific ligands. *Proc. Natl. Acad. Sci. U. S. A.* 74:5031–5035.
- Bykovskaya, S. N., A. N. Rylenko, M. O. Rauschenbach, and A. F. Bykovsky. 1978. Ultrastructural alteration of cytolytic T-lymphocytes following their interaction with target cells. I. Hypertrophy and change of orientation of the Golgi apparatus. *Cell Immunol.* 40:164–174.
- Bykovskaya, S. N., A. N. Rylenko, M. O. Rauschenbach, and A. F. Bykovsky. 1978. II. Morphogenesis of secretory granules and intracellular vacuoles. *Cell Immunol.* 40:175–185.
- Canty, T. G., and J. R. Wunderlich. 1970. Quantitative *in vitro* assay of cytotoxic cellular immunity. *J. Natl. Cancer Inst.* 45:761–772.
- Cerottini, J.-C., and K. T. Brunner. 1974. Cell-mediated cytotoxicity, allograft rejection and tumor immunity. *Adv. Immunol.* 18:67–132.
- Fishelson, Z., and G. Berke. 1978. T-lymphocyte-mediated cytotoxicity: dissociation of the binding and lytic mechanisms of the effector cell. *J. Immunol.* 120:1121–1126.
- Flanagan, J., and G. L. E. Koch. 1978. Cross-linked surface Ig attaches to actin. *Nature (Lond.)*. 273:278–281.
- Geiger, B., and S. J. Singer. 1980. Association of microtubules and intermediate filaments in chicken gizzard cells as detected by double immunofluorescence. *Proc. Natl. Acad. Sci. U. S. A.* 77:4769–4773.
- Golstein, P. 1974. Sensitivity of cytotoxic T-cells to cell-mediated cytotoxicity. *Nature (Lond.)*. 252:81–83.
- Golstein, P., and E. T. Smith. 1977. Mechanism of T-cell mediated cytotoxicity: the lethal hit stage. *Contemp. Top. Immunobiol.* 7:273–300.
- Henney, C. S. 1977. T-cell-mediated cytotoxicity: an overview of some current issues. *Contemp. Top. Immunobiol.* 7:245–272.
- Korn, E. D. 1978. Biochemistry of actomyosin-dependent cell motility (a review). *Proc. Natl. Acad. Sci. U. S. A.* 75:588–599.
- Malech, H. L., R. K. Root, and J. F. Gallin. 1977. Structural analysis of human neutrophil migration. Centriole, microtubule, and microfilament orientation and function during chemotaxis. *J. Cell Biol.* 75:666–693.
- Marshak-Rothstein, A., P. Fink, T. Gridley, D. H. Raulet, M. J. Bevan, and M. L. Gelfer. 1979. Properties and applications of monoclonal antibodies directed against determinants of the Thy-1 locus. *J. Immunol.* 122:2491–2497.
- Martz, E. 1977. Mechanism of specific tumor-cell lysis by alloimmune T-lymphocytes: resolution and characterization of discrete steps in the cellular interaction. *Contemp. Top. Immunobiol.* 7:301–361.
- de Petris, S., and M. C. Raff. 1972. Distribution of immunoglobulin on the surface of mouse lymphoid cells as determined by immunoferritin electron microscopy. Antibody-induced, temperature-dependent redistribution and its implications for membrane structure. *Eur. J. Immunol.* 2:523–535.
- Pollard, T. D., and R. P. Weibing. 1974. Actin and myosin and cell movement. *CRC Crit. Rev. Biochem.* 2:1–65.
- Rosen, D., Z. Fishelson, and G. Berke. 1981. The role of cytotoxic T-lymphocyte projections in target cell lysis. *Transplant. Proc.* 13:1073–1078.
- Ryser, J. E., E. Rungger-Brandle, C. Chaponnier, G. Gabbiani, and P. Vassalli. 1982. The area of attachment of cytotoxic T lymphocytes to their target cells shows high motility and polarization of actin, but not myosin. *J. Immunol.* 128:1159–1162.
- Sanderson, C. J. 1981. The mechanism of lymphocyte-mediated cytotoxicity. *Biol. Rev. Camb. Philos. Soc.* 56:153–197.
- Sanderson, C. J., and A. M. Glauert. 1979. The mechanism of T-cell mediated cytotoxicity. VI. T-cell projections and their role in target cell killing. *Immunology*. 36:119–129.
- Schreiner, G. F., and E. R. Unanue. 1976. Membrane and cytoplasmic changes in B lymphocytes induced by ligand and immunoglobulin interaction. *Adv. Immunol.* 24:137–165.
- Stackpole, C. W., J. B. Jacobson, and M. P. Lardis. 1974. Two distinct types of capping of surface receptors on mouse lymphoid cells. *Nature (Lond.)*. 248:232–234.
- Tilney, L. G., and J. R. Gibbons. 1969. Microtubules in the formation and development of the primary mesenchyme in *Arbacia punctulata*. II. An experimental analysis of their role in development and maintenance of cell shape. *J. Cell Biol.* 41:227–250.
- Toh, B. H., and G. C. Hard. 1977. Actin coprecipitates with concanavalin A receptors. *Nature (Lond.)*. 269:695–697.
- Trelstad, R. L. 1970. The Golgi apparatus in chick corneal epithelium: changes in intracellular position during development. *J. Cell Biol.* 45:34–42.
- Trinkaus, J. P. 1976. On the mechanism of metazoan cell movement. In *The Cell Surface in Animal Embryogenesis and Development*. G. Poste and G. L. Nicolson, editors. Elsevier/North Holland Biomedical Press, Amsterdam. 225–329.
- Unanue, E. R., W. D. Perkins, and M. J. Karnovsky. 1972. Ligand-induced movement of lymphocyte membrane macromolecules. I. Analysis of immunofluorescence and ultrastructural radioautography. *J. Exp. Med.* 136:885–906.
- Yahara, I., and G. M. Edelman. 1972. Restriction of the mobility of lymphocyte immunoglobulin receptors by concanavalin A. *Proc. Natl. Acad. Sci. U. S. A.* 69:608–612.
- Yahara, I., and G. M. Edelman. 1975. Modulation of lymphocyte receptor mobility by concanavalin A and colchicine. *Ann. N. Y. Acad. Sci.* 253:455–469.
- Yahara, I., and F. Kakimoto-Sameshima. 1978. Microtubule organization of lymphocytes and its modulation by patch and cap formation. *Cell*. 15:251–259.
- Zagury, D., J. Bernard, N. Thiernesse, M. Feldman, and G. Berke. 1975. Isolation and characterization of individual functionally reactive cytotoxic T lymphocytes: conjugation, killing and recycling at the single cell level. *Eur. J. Immunol.* 5:818–822.
- Zagury, D., J. Bernard, P. Jeannesson, N. Thiernesse, and J.-C. Cerottini. 1979. Studies on the mechanism of T cell-mediated lysis at the single effector cell level. I. Kinetic analysis of lethal hits and target cell lysis in multicellular conjugates. *J. Immunol.* 123:1604–1609.
- Zinkernagel, R. M., and P. C. Doherty. 1979. MHC-restricted cytotoxic T cells. Studies on the biological role of polymorphic major transplantation antigens determining T-cell restriction specificity, function and responsiveness. *Adv. Immunol.* 27:51–177.