

SUPPLEMENTARY DATA

The DIA-AID 1 Writing Group

Domingo Acosta-Delgado, University Hospital Virgen del Rocío, Seville, SP; Riitta Liisa Antikainen, Tutkimusyksikkö, Oulu, FI; Khajag Azejian, Tikkurilan Terveyskeskus (Tikkurila Health Center), Vantaa, FI; Ilaria Barchetta, Università "La Sapienza" di Roma, Rome, IT; Marco Giorgio Baroni, Diabetes Unit, Azienda Ospedaliero-Universitaria di Cagliari Policlinico Universitario, Monserrato, IT; Pierre Yves Benhamou, CHU de Grenoble, Grenoble, FR.; Bernhard Böhm, Universitätsklinikum Ulm, Ulm, DE; Susan Brown, Donald Gordon Medical Center, Johannesburg, SA; Maria Gisella Cavallo, Università "La Sapienza" di Roma, Rome, IT; Franco Cavalot, Internal Medicine and Metabolic Disease Unit Department of Clinical and Biological Sciences of the Turin University San Luigi Gonzaga Hospital, Orbassano, IT; Massimo Chirio, Internal Medicine and Metabolic Disease Unit Department of Clinical and Biological Sciences of the Turin University San Luigi Gonzaga Hospital, Orbassano, IT; Efisio Cossu, Diabetes Unit, Azienda Ospedaliero-Universitaria, di Cagliari Policlinico Universitario, Monserrato, IT; José M. Cubero, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma, Barcelona, SP; Thomas Danne, Diabetes Centre for Children and Adolescents, Kinderkrankenhaus auf der Bult., Hannover, DE; Nicolin Datz, Diabetes Centre for Children and Adolescents, Kinderkrankenhaus auf der Bult., Hannover, DE; Elvira Fioriti, University Campus Bio-Medico, Rome, IT; Stefano Genovese, Istituto Clinico Humanitas, Rozzano, IT; José-Miguel González-Clemente, Hospital de Sabadell, SP; Chiara Guglielmi, University Campus Bio-Medico, Rome, IT; Vesa Ilvesmäki, Päijät-Hämeen Keskussairaala, Lahti, FI; Vuokko Kallioniemi, Diabetestutkimus, Tampere, FI; Shadi Kyanvash, University Campus Bio-Medico, Rome, IT; Olga Kordonouri, Diabetes Centre for Children and Adolescents, Kinderkrankenhaus auf der Bult., Hannover, DE; Sudhesh Kumar, Walsgrave Hospital, Coventry, UK; Alberto de Leiva, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma, Barcelona, SP; Dan Lender, Diabetes Unit, Hadassah University Hospital, Jerusalem, Israel; Richard David Graham Leslie, St. Bartholomew's Hospital, London, UK; Esther Lieberman, Soroka University Medical Center; Beer Sheva, ISR; Iñaki Llorente, Hospital Universitario Nuestra, Tenerife, SP; Katerina Loykova, Faculty Hospital Olomouc, CZ; Andrew Macleod, Hummingbird Centre, Royal Shrewsbury Hospital, Shrewsbury UK; Jussi Malmi, Pohjois-Karjala Projektin Tutkimussäätiö, Joensuu, FI.; Erika Marquardt, Diabetes Centre for Children and Adolescents, Kinderkrankenhaus auf der Bult., Hannover, DE; Ofri Mosenzon, Diabetes Unit, Hadassah University Hospital, Jerusalem, Israel; Terezie Pelikánová, IKEM/Diabetes Centre, Prague, CZ; Pavlina Pithova, Faculty Hospital Motol, Prague, CZ; Paolo Poy, Internal Medicine and Metabolic Disease Unit Department of Clinical and Biological Sciences of the Turin University San Luigi Gonzaga Hospital, Orbassano, IT; Sotirios A. Raptis, Research Institute and Diabetes Centre, Athens, GR; Leslie Ivan "Mac" Robertson, 102 Parklands Medical Centre, Durban, SA; Michel Rodier, CHU de Nîmes/ Hôpital Caremeau, Nîmes, FR; Jaroslav Rybka, Regional Hospital Bata a.s., Zlín, CZ; Kerstin Schnell, Diabetes Centre for Children and Adolescents, Kinderkrankenhaus auf der Bult., Hannover, DE; Zdenek Sumnik, Faculty Hospital Motol, Prague, CZ; Nikolaos Tentolouris, LAIKO Hospital, Athens, GR; Charles Thivolet, Hopital Edouard Herriot, Lyon, FR; Mariella Trovati, Internal Medicine and Metabolic Disease Unit Department of Clinical and Biological Sciences of the Turin University San Luigi Gonzaga Hospital, Orbassano, IT; Christian Tsioli, Diabetes Centre for Children and Adolescents, Kinderkrankenhaus auf der Bult., Hannover, DE; Dario Tuccinardi, University Campus Bio-Medico, Rome, IT; Jaakko Tuomilehto, University of Helsinki, Helsinki, FI.; Jan Vavrinec, Faculty Hospital Motol, Prague, CZ; Bernard Vialettes, Hopital La Timone, Marseille, FR; Jeffery Robert Wing, Donald Gordon Medical Center, Johannesburg, SA; Holger Oswald Wellmann, Helderberg Clinical Trials Unit, Cape Town, SA; and Donald Whitelaw, Bradford Royal Infirmary, Bradford, UK.

SUPPLEMENTARY DATA

Supplementary Table 1. Parameters of Patients Screened for Eligibility.

Parameter	Average ± SD (minimum - maximum)
Gender	457 Males / 222 Females
Age (years)	27.4 ± 7.92 (15 - 47)
Time from diagnosis (months)	1.87 ± 1.27 (0 – 8.57)
BMI (kg/m ²)	22.9 ± 3.43 (16 – 39.5)
Fasting C-peptide (nmol/l)	0.46 ± 0.30 (0.07 – 2.46)
HbA1c (%)	7.52 ± 1.88 (4 - 17)
Autoantibodies (% positive)	IA2A 51.2%, IAA 65.8%, GADA 75.4%

Supplementary Table 2. Inclusion / Exclusion Criteria that were Violated by Patients Excluded from ITT to Form the mITT Population Prior to Unmasking.

Inclusion / Exclusion Criteria Violated	Number of Subjects
No affirmation of T1D	12
Time from Diagnosis (not newly diagnosed)	5
Age at Randomization (either <16 or >45)	7
Ketonuria & proteinuria at visit 1; Use of Metformin	1
Repeated tests of ketonuria, proteinuria, hematuria	2
Ketones in urine and weight loss > 10%	1
No insulin taken for at least 2 months after diagnosis	3
Underlying disease	1
Received both allocations (treatment group cannot be determined for efficacy evaluation)	2

Calculations Used for Data Analysis

Change from baseline in AUC = $AUC_{24\text{ months}} - AUC_{\text{baseline}}$

Relative treatment effect = $(1 - \text{LS mean}_{\text{DiaPep277}}^{\text{®}} / \text{LS mean}_{\text{Placebo}}) \times 100$.

Rate of hypoglycemic events/month was calculated as follows (based on 30 days per month):

Rate of Events/month at Baseline = $\text{Number of Events} / \{[(\text{Month 3} - \text{Baseline}) + 1] / 30\}$; Rate of

Events/month at Study End = $\text{Number of Events} / \{[(\text{Month 24} - \text{Month 3}) + 1] / 30\}$.

Change of Hypoglycemic Event Rate = $(\text{Hypoglycemic event rate at Month 24} - \text{Hypoglycemic event rate at Month 3})$.

SUPPLEMENTARY DATA

Supplementary Table 3. Hypoglycemia Classification.

Severity	Symptoms	Plasma Glucose	Action Taken	Classification
Subject able to initiate self-treatment if necessary	Symptoms suggestive hypoglycemia of	< 70 mg/dL (3.9 mmol/L)	Enter in Glycemia Study Diary eCRF including the actions taken	Symptomatic Hypoglycemia*
	Symptoms suggestive hypoglycemia of	≥70 mg/dL (3.9 mmol/L)	Enter in Glycemia Study Diary eCRF including the actions taken	Relative Hypoglycemia*
	Symptoms suggestive hypoglycemia of	Not taken	Enter in Glycemia Study Diary eCRF including the actions taken	Probable symptomatic Hypoglycemia*
	Asymptomatic	< 70 mg/dL (3.9 mmol/L)	Enter in Glycemia Study Diary eCRF including the actions taken	Asymptomatic Hypoglycemia*
Subject is unable to initiate self-treatment and requires assistance of another person or hospitalization.	Symptoms suggestive hypoglycemia of	< 70 mg/dL (3.9 mmol/L)	Enter in Glycemia Study Diary eCRF Report as SAE if criteria are fulfilled including the actions taken	Severe Hypoglycemia* SAE, if applicable
	Symptoms suggestive hypoglycemia of	Clinical manifestation reversed by administration of oral carbohydrate, subcutaneous glucagon or intravenous glucose	Enter in Glycemia Study Diary eCRF Report as SAE if criteria are fulfilled including the actions taken	Severe Hypoglycemia* SAE, if applicable

SUPPLEMENTARY DATA

Supplementary Table 4. Reasons for Screen Failure.

	Number	Percentage
Autoantibodies negative	54	24.32
Low C-peptide	54	24.32
Withdrawal of Consent	40	18.02
Failed to comply with inclusion/exclusion criteria	21	9.46
Patient not compliant with requirements of the study	19	8.56
Aberrant lab results	10	4.50
Time from Diagnosis not applicable	8	3.60
Lost to follow-up	5	2.25
BMI out of range	4	1.80
Age	2	0.90
Unexplained weight loss	2	0.90
Allergy to penicillin	1	0.45
Worsening or complication of the underlying disease	1	0.45
Hyperglycemic Event	1	0.45
Total	222	100

Supplementary Table 5. Drop-out Patients: Breakdown by Reason and Allocation.

Reasons for not Completing the Study	Total	Breakdown by Allocation	
		DiaPep277 [®]	Placebo
Withdrawal of Consent	36	17	19
Lost to follow-up	19	6	13
Patient not compliant with requirements of the study	10	6	4
Adverse Event	9	6	3
HbA1c withdrawal criteria	8	5	3
Serious Adverse Event	4	2	2
Dermal Hypersensitivity	3	0	3
Other	3	2	1
Patient's findings or conduct failed to meet protocol entry criteria	3	2	1
Patient requires use of an unacceptable medication	2	0	2
Death	1	1	0
Other: Occurrence of a concomitant disease	1	0	1
Pregnancy	1	1	0
Protocol Violation	1	1	0
Termination by sponsor	1	1	0
Total	102	50	52
Patients included in the primary efficacy endpoint analysis	35	18	17

SUPPLEMENTARY DATA

Supplementary Table 6. Exploratory Analysis, The Proportion of Patients in Partial Remission at Study End (HbA1c \leq 7% and Insulin Dose \leq 0.5 U/kg/day).

	% of Patients		p value
mITT Population			
	DiaPep277 [®] (N = 159)	Placebo (N = 167)	
HbA1c \leq 7% & Insulin Dose \leq 0.5 U/kg/day	38.4	29.3	0.085
HbA1c >7% & Insulin Dose >0.5 U/kg/day	27	37.7	0.04
PP Population			
	DiaPep277 [®] (N = 146)	Placebo (N = 162)	
HbA1c \leq 7% & Insulin Dose \leq 0.5 U/kg/day	41.8	30.2	0.035
HbA1c >7% & Insulin Dose >0.5 U/kg/day	23.3	37.7	0.006

Supplementary Table 7. Overall Summary of Treatment Emergent Adverse Events.

	DiaPep277 [®] (N=225)	Placebo (N=231)	Total (N=456)
Number of Patients with at Least One TEAE	173 (76.9%)	164 (71%)	337 (73.9%)
Number of Patients with at Least One Serious TEAE Not Considered to be Drug Related	23 (10.2%)	13 (5.6%)	36 (7.9%)
Number of Patients with at Least One Suspected Drug-Related Serious TEAE	3 (1.3%)	1 (0.4%)	4 (0.9%)
Number of Patients with at Least One Life threatening TEAE	2 (0.9%)	2 (0.9%)	4 (0.9%)
Number of Patients with at Least One Severe TEAE	20 (8.9%)	13 (5.6%)	33 (7.2%)
Number of Patients with at Least One Suspected Drug-Related AE	31 (13.8%)	34 (14.7%)	65 (14.3%)
Number of Subject Deaths	1 (0.4%)	1 (0.4%) ¹	2 (0.4%)

N= number of patients; TEAE= treatment emergent adverse event; AE= adverse event

¹This life-threatening event and death documented in the Placebo group did not occur in a study patient, but in the neonate of a male patient (Patient 4309-002). The partner of the patient became pregnant during the study and gave birth to twins prematurely. One of the twin babies died during the study. The life-threatening event and death was recorded in the database under the Placebo group. There were no significant differences in the TEAEs between the Diapep277[®] treated and the Placebo treated groups by Fisher's exact test.


SUPPLEMENTARY DATA

Supplementary Fig 1.


	Screening Period (1 month)	Treatment Period (21 months)									Follow-up Period (3 months)	
Visit		1	2	3	4	5	6	7	8	9	10	
		↑	↑	↑	↑	↑	↑	↑	↑	↑	↑	
Month	-1	0	1	3	6	9	12	15	18	21	24	
		Baseline Randomization										

↑ Application of blinded study medication

Supplementary Figure 2. Change from Baseline in Glucagon-Stimulated Peak C_{max} Secretion.
A



B



Mean changes (\pm SE of means) in maximum C-peptide concentration (C_{max}) as determined using the 20 min glucagon stimulation test in (A) the mITT population and (B) the PP population. * $P < 0.05$.