
Beauty is in the Genes of the Beholder

A Golden Tribute to a Golden Anniversary*



By David Harel
Department of Computer
Science and Applied
Mathematics, The
Weizmann Institute of
Science, Rehovot, Israel




Ron Unger Faculty of Life Sciences, Bar-Ilan University, Ramat Gan, Israel




Joel L. Sussman
Department of Structural
Biology, The Weizmann
Institute of Science,
Rehovot, Israel

*With apologies to Watson and Crick










Fig. 1. Golden ratio in DNA

this claim with the use of the Golden Ratio. a concept going back to ancient times. The Golden Ratio, denoted by φ, is an irrational number, like π , which expresses a fundamental ratio that is almost as common as π and has the habit of appearing when least expected [2]. The geometrical meaning of ϕ is seen from the line segments of Fig. 2. The horizontal length of the rectangle has been divided into two parts, A and B, such that the ratio of B to A is the same as that of A to A+B, the length of the entire line. In each case the ratio can easily be shown to be given by:

$$\phi = \frac{1 + \sqrt{5}}{2}$$

or approximately 1.61803398.

There is little doubt that the Ancient Greeks were familiar with the Golden Ratio and it was used by some of their architects and sculptors, particularly in the structure of the Parthenon (see Fig. 3a). In fact, the name ϕ was originally given by the American mathematician Mark Barr in the early 1900s in memory of the famous Greek sculptor Phidias, who often used the Golden Ratio in

The number ϕ appears in Renaissance art, including works of da Vinci and others, and is rumored to be the basis of the segmentation of a well-built human body; first finger joint to second, second to both, hand to lower arm, lower arm to hand+lower arm, and so on, culminating in navel upwards to navel downwards, and navel downwards to whole body. This particular observation strengthens one's feeling that the proportions of DNA should similarly be based on ϕ .

That the structure of a double helix is in itself one of immense beauty goes without saying; we concentrate here on the local dimensions of the type-B double helix, which appears to be the most common form of DNA in chromatin3. It is easy to conceive of a double helix of uncomely proportions, but it is difficult to imagine that one which was unpleasant to behold would be chosen to




Fig. 2 Schematic illustration of the Golden Ratio: (A+B)/A = A/B, so that both the external and internal rectangles are identically proportioned.





Fig.3 The Golden Ratio in art and nature. (a) The Parthenon: manmade beauty; (b) DNA: natural beauty.



Section through a Nautilus shell.

If a golden rectangle is subdivided by a line which forms a square out of one end, the remainder is another. smaller golden rectangle. Repeated subdivision results in a series of squares and golden rectangles which spiral into the center of the pattern.



carry the genetic code. Given that DNA is to be structured as a right-handed double helix, there are three crucial dimensions that essentially determine its final form: the external width (diameter) of the double helix, the length of its period (i.e. the pitch, or the height of a "slice" thereof), and the vertical offset of one helix from the other, which forms the minor groove.

The novel feature of our interpretation of Watson and Crick's structure is the manner in which these three basic dimensions are held together by the rule of the Golden Ratio. Specifically we have found that the ratio of the height to the width and the ratio of the width of the offset are always very close to o (see Figs. 1 and 3). In particularly, since the ratio between height and offset is responsible for the unequal sizes of the major and minor grooves of B-DNA, we might say that the ratio between the grooves themselves is golden too.

Although the crystal structure of a B-DNA dodecamer4 has been refined to high resolution, it is too short and irregular to accurately estimate the pitch, diameter and offset of the DNA molecule as a whole. However, the most recent and best refined X-ray fibre studies of long chains of B-DNA5 (re-refined by Arnott, S. and Chandrasekaran, R., personal communication 1928) yield ratios of 1.6031 and 1.538, respectively. or approximately 1 and 5% less than φ. Moreover, both ratios seem to converge to o as time proceeds (Fig. 4). In contrast, in the less frequently occurring forms of DNA, i.e. A-DNA and Z-DNA, the ratios are further from φ, as shown in Table I.

Currently available X-ray data on DNA are insufficient for a rigorous test of our interpretation of the structure. As far as we can tell, however, our interpretation is smoothly compatible with the experimental data (and is getting better with time). In fact, just 7 years after the Watson-Crick model was proposed.

Table I. Models of DNA structures and their fit to the Golden Ratio

Model	Height (Å) (pitch)	Width (Å) ^a (diameter)	Offset (Å)b	Height/ width	Width/ offset
A-DNA⁵,C	25.6	18.9	19.1	1.85	0.99
B-DNA⁵, C	33.8	21.1	13.7	1.60	1.54
Z-I DNA9	44.6	17.5	15.9	2.55	1.10
Z-II DNA9	44.6	18.3	10.9	2.44	1.68

^aWidth determined by the atom most distant from the helix axis: A-DNA 05', B-DNA 02P, Z-I DNA C8(G), Z-II DNA 02P (C).

Offset determined by vertical separation between the two helices of the double helix based on the locus of phosphorus atoms.

Re-refined by Arnott, S. and Chandrasekaran, R., personal communication (1982).




Fig. 4. Beauty improves in time. The coordinates for the 4 DNA structures are taken from Refs 5 (re-refined by Arnott, S and Chandrasekaran, R., personal communication, 1982) and 6-8.

Langridge et al.10 built a DNA model with ratios of height/width = 1.63 and width/offset = 1.51. But it must be regarded as unproven until it has been checked against more exact results. Our 99% and 95% fits, in other words, are only the starting point...

It has not escaped our notice that the specific interpretation we have postulated immediately suggests a possible motto for the copying mechanism of the genetic material:

Beauty is in the genes of the beholder

Joel L. Sussman is incumbent of the Morton and Gladys Pickman Professorial Chair in Structural Biology and David Harel is incumbent of the William Sussman Professorial Chair in Mathematics.

This article first appeared under the title "Beauty is in the Genes of the Beholder", in Trends Biochem. Sci., Vol 11, (1986), 155-156. Reproduced with kind permission of Reed Elsevier.

References

- 1 Watson, J.D. and Crick, F.H.C. (1953) Nature 171, 737-738
- 2 Gardner, M. (1959) Sci. Am. 200, 128-134
- 3 Richmond, T.J., Finch, J.T., Rushton, B., Rhodes, D. and Klug, A. (1984) Nature 311, 532-537
- 4 Wing, R., Drew, H., Takano, T., Broka, C., Tanaka, S., Itakura, K. and Dickerson, R.E. (1980) Nature 287, 755-758
- 5 Arnott, S., Chandrasekaran, R., Birdsall, D.L., Leslie, A.G.W. and Ratliff, R.L. (1980) Nature 283, 743-745
- 6 Crick, F.H.C. and Watson, J.D. (1954) Proc. Roy. Soc. A223,
- 7 Arnott, S., Dover, S.D. and Wonacott, A.J. (1969) Acta Crystallogr. B25, 2192-2206
- 8 Arnott, S. and Hukins, D.W.L. (1972) Biochem. Biophys. Res. Comm. 47, 1504-1509
- 9 Wang, A.H-J., Quigley, G.J., Kolpak, F.J., Crawford, J.L., van Boom, J.H., van der Marel, G. and Rich, A. (1979) Nature 282,
- 10 Langridge, R., Marvin, D.A., Seeds, W.E., Wilson, U.R., Hooper, C.W., Wilkins, M.U.F. and Hamilton, L.D. (1960) J. Mol. Biol. 2, 38-64.