

# Introduction to Systems Neuroscience

Methodologies used to study brain systems


Ehud Ahissar

Department of Neurobiology

# Introduction


- In natural sciences, unlike in mathematics, there is ***no set of pre-defined axioms*** from which other definitions can be derived and proved.
- What can be proven in natural sciences are only ***consistencies***; hypotheses are considered valid as long as they are consistent with actual observations
  - *e.g., gravity theory is considered valid as long as the movement patterns it entails are consistent with actual movements of bodies;*
  - *“gravity” was never “observed” by itself.*
- The methods are thus developed to test consistencies of observations with hypotheses

# Introduction



# Introduction

## The scientific method



# Research approaches

- **Observational:** we collect enough evidence to compose a hypothesis
- **Correlational:** we compare 2 variables, the values of which have been collected without direct intervention. No causal relationships can be directly concluded.
- **Experimental (Interventional):** an “independent” variable is systematically manipulated and the effects of this on a “dependent” variable are measured. Considered as the best approach for revealing causal relationships, although...
  - It assumes that the manipulator (e.g., the experimenter) is external to the studied system
  - A problematic assumption at the frontiers of science today:
    - Quantum mechanics
    - neuroscience

# Introduction

## Examples of general guiding assumptions

*(Practically not challenged by anyone in the field)*

- All mental functions are carried out in the brain
- Brain processes obey the rules of physics

## Examples of common working hypotheses

*(often specific to research groups and research topic)*

- “The basic component of processing is a single neuron”
- “Network processing is mediated by neurotransmitters, brain states are controlled by neuromodulators”
- “Memories are stored in synapses”

# Introduction



## Major difficulties

- Complexity
    - many levels (ions, ... , neurons, ..., systems)
    - many components (e.g.,  $10^{11}$  neurons,  $10^{15}$  synapses)
    - many variables (physical, chemical, mechanical, electrical, physiological, psychological)
  - Small sizes (neurons  $\sim 10 \mu\text{m}$ , synapse  $< 1 \mu\text{m}$ )
  - Closed loops
  - Self organization
  - plasticity
- } Cannot repeat an experiment twice

# Introduction

## Major difficulties – Methods to address them

- Complexity

- many levels (ions, ... , ne  systems) ...
- many components (e.g.  ions,  $10^{15}$  synapses)
- many variables (physical, chemical, mechanical, electrical, physiological, psychological)

- Small sizes (neurons

Small tools, magnifications, genetic tools

- Closed loops

Opening the loops by anesthesia, flashed stimuli, cuts

- Self organization

- plasticity

Cannot repeat an experiment twice

Using statistics

# Introduction

## About this lecture

- Review of popular methods for experiments in systems neuroscience
- Focus on resolutions and potential power. Less on limitations
- Not all methods will be covered. You are invited to complete the picture through the web or text books
- Slide numbers – appear in MOST of the slides, not in all...

# Methods table

## Measuring neural activity

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>behavior</b>	brain	brain	$> 10^{11}$	station	1 ms	10 ms	10	
<b>2DG, c-fos</b>	10 $\mu\text{m}$	10 $\mu\text{m}$	1		NA	30 min	$> 10^6$	
<b>fMRI</b>	$> 1 \text{ mm}$	$> 200 \mu\text{m}$	$> 10^5$		1 s	$> 1 \text{ s}$	1000	10 ms
<b>EEG</b>	10 mm	100 mm	$> 10^9$	station	$< 1 \text{ ms}$	50 ms	50	1 ms
<b>MEG</b>	10 mm	100 mm	$> 10^9$		1 ms	50 ms	50	
<b>ECoG</b>	10 mm	1-10 mm	$> 10^{4-6}$		$< 1 \text{ ms}$	10 ms	10	
<b>Intr. Signal</b>	10 $\mu\text{m}$	200 $\mu\text{m}$	$> 1000$		$< 1 \text{ ms}$	1 s	1000	10 ms
<b>VSD Imaging</b>	10 $\mu\text{m}$	30 $\mu\text{m}$	$< 100$		$< 1 \text{ ms}$	10 ms	$< 1$	
<b><math>\mu\text{Elec}</math></b>	50 $\mu\text{m}$	10 $\mu\text{m}$	1	10 $\mu\text{m}$	$< 1 \text{ ms}$	1 ms	1	
<b><math>\mu\text{Dialysis}</math></b>	$< 1 \text{ mm}$	100 $\mu\text{m}$	$> 10^4$		$> 1 \text{ min}$	100 ms	$> 10^6$	
<b>Intracell elec</b>	10 $\mu\text{m}$	10 $\mu\text{m}$	$< 1$		$< 1 \text{ ms}$	$< 1 \text{ ms}$	$< 1$	
<b>Ca imaging</b>	1 $\mu\text{m}$	1 $\mu\text{m}$	$< 1$		1 ms	$> 100 \text{ ms}$	$> 100$	

# Methods table

Stimulating / perturbing neural activity

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>sensory</b>	modality	modality	$> 10^9$	station	$< 1$ ms	$< 1$ ms	$< 1$	
<b>TMS</b>	100 mm	10 mm	$> 10^9$	station	$< 1$ ms	100 ms	100	
<b><math>\mu</math>Stim</b>	$< 10 \mu\text{m}$	$> 100 \mu\text{m}$	$> 50$		$< 1$ ms	10 ms	10	
<b><math>\mu</math>Pharmac</b>	$< 10 \mu\text{m}$	$> 100 \mu\text{m}$	$> 50$		1 ms	$> 10$ s	10	
<b>single cell</b>	$< 10 \mu\text{m}$	$< 10 \mu\text{m}$	1		$< 1$ ms	$< 1$ ms	$< 1$	
<b>sub-cell</b>	$> 100 \mu\text{m}$	$< 1 \mu\text{m}$	$> 50$		$< 1$ ms	$< 1$ ms	$< 1$	

# Methods table

## Measuring structure (anatomy)

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>cell density</b>								
<b>receptor density</b>								
<b>transmitter density</b>								
<b>tract tracing</b>								
<b>single-cell</b>								

# Methods table

## Manipulating structure

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>Neuropsychology</b>		> 10 mm	> 10 <sup>7</sup>			months		
<b>lesions</b>	> 100 μm	> 100 μm	> 50		> 1 s	> 1 min		

# Measuring neural activity

## Behavior

- Present sensory stimuli
- Measure response accuracy, threshold, speed
- Infer related neural activity

Example: roughness perception

- Present boards with varying roughness
- Ask to estimate roughness (say [1:10])
- Compute a psycho-physical curve

Examples... estimation (x-y), discrimination (dx-dy)

- Assessing threshold: the staircase paradigm
- Perceptual speed: measure reaction time

Heuristics: inferring related neuronal activity by prior knowledge

e.g., figuring station resolution by known functional anatomy

Example – tactile regions versus visual or olfactory, cortex versus brainstem, ...

## Psychophysical curve

(Reported roughness)

Psycho variable



Physical variable

(Physical density)

# Methods table

## Measuring neural activity

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
behavior	brain	brain	$> 10^{11}$	station	1 ms	10 ms	10	
2DG, c-fos	10 $\mu\text{m}$	10 $\mu\text{m}$	1		NA	30 min	$> 10^6$	
fMRI	$> 1 \text{ mm}$	$> 200 \mu\text{m}$	$> 10^5$		1 s	$> 1 \text{ s}$	1000	10 ms
EEG	10 mm	100 mm	$> 10^9$	station	$< 1 \text{ ms}$	50 ms	50	1 ms
MEG	10 mm	100 mm	$> 10^9$		1 ms	50 ms	50	
ECoG	10 mm	1-10 mm	$> 10^{4-6}$		$< 1 \text{ ms}$	10 ms	10	
Intr. Signal	10 $\mu\text{m}$	200 $\mu\text{m}$	$> 1000$		$< 1 \text{ ms}$	1 s	1000	10 ms
VSD Imaging	10 $\mu\text{m}$	30 $\mu\text{m}$	$< 100$		$< 1 \text{ ms}$	10 ms	$< 1$	
$\mu\text{Elec}$	50 $\mu\text{m}$	10 $\mu\text{m}$	1	10 $\mu\text{m}$	$< 1 \text{ ms}$	1 ms	1	
$\mu\text{Dialysis}$	$< 1 \text{ mm}$	100 $\mu\text{m}$	$> 10^4$		$> 1 \text{ min}$	100 ms	$> 10^6$	
Intracell elec	10 $\mu\text{m}$	10 $\mu\text{m}$	$< 1$		$< 1 \text{ ms}$	$< 1 \text{ ms}$	$< 1$	
Ca imaging	1 $\mu\text{m}$	1 $\mu\text{m}$	$< 1$		1 ms	$> 100 \text{ ms}$	$> 100$	

# Methods table


## Measuring neural activity

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
behavior	brain	brain	$> 10^{11}$	station	1 ms	10 ms	10	
2DG, c-fos	10 $\mu\text{m}$	10 $\mu\text{m}$	1		NA	30 min	$> 10^6$	
fMRI	$> 1 \text{ mm}$	$> 200 \mu\text{m}$	$> 10^5$		1 s	$> 1 \text{ s}$	1000	10 ms
EEG	10 mm	100 mm	$> 10^9$	station	$< 1 \text{ ms}$	50 ms	50	1 ms
MEG	10 mm	100 mm	$> 10^9$		1 ms	50 ms	50	
ECoG	10 mm	1-10 mm	$> 10^{4-6}$		$< 1 \text{ ms}$	10 ms	10	
Intr. Signal	10 $\mu\text{m}$	200 $\mu\text{m}$	$> 1000$		$< 1 \text{ ms}$	1 s	1000	10 ms
VSD Imaging	10 $\mu\text{m}$	30 $\mu\text{m}$	$< 100$		$< 1 \text{ ms}$	10 ms	$< 1$	
$\mu\text{Elec}$	50 $\mu\text{m}$	10 $\mu\text{m}$	1	10 $\mu\text{m}$	$< 1 \text{ ms}$	1 ms	1	
$\mu\text{Dialysis}$	$< 1 \text{ mm}$	100 $\mu\text{m}$	$> 10^4$		$> 1 \text{ min}$	100 ms	$> 10^6$	
Intracell elec	10 $\mu\text{m}$	10 $\mu\text{m}$	$< 1$		$< 1 \text{ ms}$	$< 1 \text{ ms}$	$< 1$	
Ca imaging	1 $\mu\text{m}$	1 $\mu\text{m}$	$< 1$		1 ms	$> 100 \text{ ms}$	$> 100$	

# MRI vs. fMRI

MRI

high  
resolution  
(1 mm)



one image

fMRI

low resolution  
(~3 mm but can be better)




many images  
(e.g., every 2 sec for 5 mins)


fMRI

Blood Oxygenation Level Dependent (BOLD)  
signal indirect measure of neural activity

↑ neural activity → ↑ blood oxygen → ↑ fMRI signal

# fMRI using BOLD






# Predicting fMRI BOLD signal in one subject from spike activity in another subject during the same movie


Patient 1 predictor

n = 6



Correlation = 0.73,  $p \ll 0.001$

# Non-linear amplification revealed in fMRI signals




# Methods table

## Measuring neural activity

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
behavior	brain	brain	$> 10^{11}$	station	1 ms	10 ms	10	
2DG, c-fos	10 $\mu\text{m}$	10 $\mu\text{m}$	1		NA	30 min	$> 10^6$	
fMRI	$> 1 \text{ mm}$	$> 200 \mu\text{m}$	$> 10^5$		1 s	$> 1 \text{ s}$	1000	10 ms
EEG	10 mm	100 mm	$> 10^9$	station	$< 1 \text{ ms}$	50 ms	50	1 ms
MEG	10 mm	100 mm	$> 10^9$		1 ms	50 ms	50	
ECoG	10 mm	1-10 mm	$> 10^{4-6}$		$< 1 \text{ ms}$	10 ms	10	
Intr. Signal	10 $\mu\text{m}$	200 $\mu\text{m}$	$> 1000$		$< 1 \text{ ms}$	1 s	1000	10 ms
VSD Imaging	10 $\mu\text{m}$	30 $\mu\text{m}$	$< 100$		$< 1 \text{ ms}$	10 ms	$< 1$	
$\mu\text{Elec}$	50 $\mu\text{m}$	10 $\mu\text{m}$	1	10 $\mu\text{m}$	$< 1 \text{ ms}$	1 ms	1	
$\mu\text{Dialysis}$	$< 1 \text{ mm}$	100 $\mu\text{m}$	$> 10^4$		$> 1 \text{ min}$	100 ms	$> 10^6$	
Intracell elec	10 $\mu\text{m}$	10 $\mu\text{m}$	$< 1$		$< 1 \text{ ms}$	$< 1 \text{ ms}$	$< 1$	
Ca imaging	1 $\mu\text{m}$	1 $\mu\text{m}$	$< 1$		1 ms	$> 100 \text{ ms}$	$> 100$	


# EEG recording



Epileptic spike and wave discharges monitored with EEG.

# EEG recording

## Dipole Formation in EEG Potentials




From: DeFelipe and Farinas, 1992

# EEG recording

## PRINCIPLES of VOLUME CONDUCTION


The amplitude of spikes strongly decay, and therefore, they do not summate to levels that is detectable by electrodes not located in their immediate neighborhood ( $30\text{-}100\mu$ ).



# EEG recording

EEG signals reflect synchronous waves of dendritic activities

Slow signals – note the distinction between device and signal resolutions



# Methods table

## Measuring neural activity


	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
behavior	brain	brain	$> 10^{11}$	station	1 ms	10 ms	10	
2DG, c-fos	10 $\mu\text{m}$	10 $\mu\text{m}$	1		NA	30 min	$> 10^6$	
fMRI	$> 1 \text{ mm}$	$> 200 \mu\text{m}$	$> 10^5$		1 s	$> 1 \text{ s}$	1000	10 ms
EEG	10 mm	100 mm	$> 10^9$	station	$< 1 \text{ ms}$	50 ms	50	1 ms
MEG	10 mm	100 mm	$> 10^9$		1 ms	50 ms	50	
ECoG	10 mm	1-10 mm	$> 10^{4-6}$		$< 1 \text{ ms}$	10 ms	10	
Intr. Signal	10 $\mu\text{m}$	200 $\mu\text{m}$	$> 1000$		$< 1 \text{ ms}$	1 s	1000	10 ms
VSD Imaging	10 $\mu\text{m}$	30 $\mu\text{m}$	$< 100$		$< 1 \text{ ms}$	10 ms	$< 1$	
$\mu\text{Elec}$	50 $\mu\text{m}$	10 $\mu\text{m}$	1	10 $\mu\text{m}$	$< 1 \text{ ms}$	1 ms	1	
$\mu\text{Dialysis}$	$< 1 \text{ mm}$	100 $\mu\text{m}$	$> 10^4$		$> 1 \text{ min}$	100 ms	$> 10^6$	
Intracell elec	10 $\mu\text{m}$	10 $\mu\text{m}$	$< 1$		$< 1 \text{ ms}$	$< 1 \text{ ms}$	$< 1$	
Ca imaging	1 $\mu\text{m}$	1 $\mu\text{m}$	$< 1$		1 ms	$> 100 \text{ ms}$	$> 100$	

# EEG recording



Epileptic spike and wave discharges monitored with EEG.

# EEG & brain states



# EEG recording

## ERP – Event Related Potentials

When an event is repeated tens or hundreds of times, and the time-related EEG signals are averaged, the resulting signal is considered as the average potential evoked by the event


# EEG recording

## ERP – Event Related Potentials




Left ear

heuristics



Right ear



Time (ms)



# Methods table

## Measuring neural activity

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>behavior</b>	brain	brain	$> 10^{11}$	station	1 ms	10 ms	10	
<b>2DG, c-fos</b>	10 $\mu\text{m}$	10 $\mu\text{m}$	1		NA	30 min	$> 10^6$	
<b>fMRI</b>	$> 1 \text{ mm}$	$> 200 \mu\text{m}$	$> 10^5$		1 s	$> 1 \text{ s}$	1000	10 ms
<b>EEG</b>	10 mm	100 mm	$> 10^9$	station	$< 1 \text{ ms}$	50 ms	50	1 ms
<b>MEG</b>	10 mm	100 mm	$> 10^9$		1 ms	50 ms	50	
<b>ECoG</b>	10 mm	1-10 mm	$> 10^{4-6}$		$< 1 \text{ ms}$	10 ms	10	
<b>Intr. Signal</b>	10 $\mu\text{m}$	200 $\mu\text{m}$	$> 1000$		$< 1 \text{ ms}$	1 s	1000	10 ms
<b>VSD Imaging</b>	10 $\mu\text{m}$	30 $\mu\text{m}$	$< 100$		$< 1 \text{ ms}$	10 ms	$< 1$	
<b><math>\mu\text{Elec}</math></b>	50 $\mu\text{m}$	10 $\mu\text{m}$	1	10 $\mu\text{m}$	$< 1 \text{ ms}$	1 ms	1	
<b><math>\mu\text{Dialysis}</math></b>	$< 1 \text{ mm}$	100 $\mu\text{m}$	$> 10^4$		$> 1 \text{ min}$	100 ms	$> 10^6$	
<b>Intracell elec</b>	10 $\mu\text{m}$	10 $\mu\text{m}$	$< 1$		$< 1 \text{ ms}$	$< 1 \text{ ms}$	$< 1$	
<b>Ca imaging</b>	1 $\mu\text{m}$	1 $\mu\text{m}$	$< 1$		1 ms	$> 100 \text{ ms}$	$> 100$	

# MEG recording

- MEG picks up magnetic fields generated by ionic currents in the brain
- Like EEG, it will show meaningful signals only for synchronized and coordinated currents.
- Like EEG, the major source of these signals are synaptic currents in cortical pyramidal neurons
- Unlike EEG, MEG is sensitive to the direction of the summed current
- As a result, comparison of MEG and EEG can increase the resolution of source localization
- So far, source localization is very limited.




# Methods table

## Measuring neural activity

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>behavior</b>	brain	brain	$> 10^{11}$	station	1 ms	10 ms	10	
<b>2DG, c-fos</b>	10 $\mu\text{m}$	10 $\mu\text{m}$	1		NA	30 min	$> 10^6$	
<b>fMRI</b>	$> 1 \text{ mm}$	$> 200 \mu\text{m}$	$> 10^5$		1 s	$> 1 \text{ s}$	1000	10 ms
<b>EEG</b>	10 mm	100 mm	$> 10^9$	station	$< 1 \text{ ms}$	50 ms	50	1 ms
<b>MEG</b>	10 mm	100 mm	$> 10^9$		1 ms	50 ms	50	
<b>ECoG</b>	10 mm	1-10 mm	$> 10^{4-6}$		$< 1 \text{ ms}$	10 ms	10	
<b>Intr. Signal</b>	10 $\mu\text{m}$	200 $\mu\text{m}$	$> 1000$		$< 1 \text{ ms}$	1 s	1000	10 ms
<b>VSD Imaging</b>	10 $\mu\text{m}$	30 $\mu\text{m}$	$< 100$		$< 1 \text{ ms}$	10 ms	$< 1$	
<b><math>\mu\text{Elec}</math></b>	50 $\mu\text{m}$	10 $\mu\text{m}$	1	10 $\mu\text{m}$	$< 1 \text{ ms}$	1 ms	1	
<b><math>\mu\text{Dialysis}</math></b>	$< 1 \text{ mm}$	100 $\mu\text{m}$	$> 10^4$		$> 1 \text{ min}$	100 ms	$> 10^6$	
<b>Intracell elec</b>	10 $\mu\text{m}$	10 $\mu\text{m}$	$< 1$		$< 1 \text{ ms}$	$< 1 \text{ ms}$	$< 1$	
<b>Ca imaging</b>	1 $\mu\text{m}$	1 $\mu\text{m}$	$< 1$		1 ms	$> 100 \text{ ms}$	$> 100$	

# ECoG recording

- To increase resolution one has to invade the brain
- The first step inside is with Electro-cortico-graphy (ECoG) – using electrodes that are placed on the surface of the brain, above or below the dura mater.
- The method is used mainly in the treatment of epilepsy, but also used to collect experimental data






# Methods table

## Measuring neural activity


	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
behavior	brain	brain	$> 10^{11}$	station	1 ms	10 ms	10	
2DG, c-fos	10 $\mu\text{m}$	10 $\mu\text{m}$	1		NA	30 min	$> 10^6$	
fMRI	$> 1 \text{ mm}$	$> 200 \mu\text{m}$	$> 10^5$		1 s	$> 1 \text{ s}$	1000	10 ms
EEG	10 mm	100 mm	$> 10^9$	station	$< 1 \text{ ms}$	50 ms	50	1 ms
MEG	10 mm	100 mm	$> 10^9$		1 ms	50 ms	50	
ECoG	10 mm	1-10 mm	$> 10^{4-6}$		$< 1 \text{ ms}$	10 ms	10	
<b>Intr. Signal</b>	10 $\mu\text{m}$	200 $\mu\text{m}$	$> 1000$		$< 1 \text{ ms}$	1 s	1000	10 ms
VSD Imaging	10 $\mu\text{m}$	30 $\mu\text{m}$	$< 100$		$< 1 \text{ ms}$	10 ms	$< 1$	
$\mu\text{Elec}$	50 $\mu\text{m}$	10 $\mu\text{m}$	1	10 $\mu\text{m}$	$< 1 \text{ ms}$	1 ms	1	
$\mu\text{Dialysis}$	$< 1 \text{ mm}$	100 $\mu\text{m}$	$> 10^4$		$> 1 \text{ min}$	100 ms	$> 10^6$	
Intracell elec	10 $\mu\text{m}$	10 $\mu\text{m}$	$< 1$		$< 1 \text{ ms}$	$< 1 \text{ ms}$	$< 1$	
Ca imaging	1 $\mu\text{m}$	1 $\mu\text{m}$	$< 1$		1 ms	$> 100 \text{ ms}$	$> 100$	

# Intrinsic signals recording


## The Cranial Window




# Left eyes was stimulated




# Right eyes was stimulated





# Left eyes was stimulated




# Right eyes was stimulated




# Ocular dominance columns



x 1000 =





1 mm 


# SOURCES OF INTRINSIC SIGNALS

- **Changes in absorption (similar to BOLD) due to:**
  - Changes in oxygenation
  - Changes in blood volume
  - Changes in blood flow
- **Changes in Light scattering due to:**

Ion movement; water movement; shrinkage or expansion of the extracellular space; transmitter release; Volume changes due to capillaries dilation

# RESOLUTION OF INTRINSIC (and BOLD) SIGNALS

Full 3-D view (no threshold) of a single WFR



**RESOLUTION depends on the threshold – Heuristic knowledge helps setting the appropriate threshold**

Barrel area  $\sim 0.15 \text{ mm}^2$ ; WFR area  $\sim 15 \text{ mm}^2$

# Methods table



## Measuring neural activity

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>behavior</b>	brain	brain	$> 10^{11}$	station	1 ms	10 ms	10	
<b>2DG, c-fos</b>	10 $\mu\text{m}$	10 $\mu\text{m}$	1		NA	30 min	$> 10^6$	
<b>fMRI</b>	$> 1 \text{ mm}$	$> 200 \mu\text{m}$	$> 10^5$		1 s	$> 1 \text{ s}$	1000	10 ms
<b>EEG</b>	10 mm	100 mm	$> 10^9$	station	$< 1 \text{ ms}$	50 ms	50	1 ms
<b>MEG</b>	10 mm	100 mm	$> 10^9$		1 ms	50 ms	50	
<b>ECoG</b>	10 mm	1-10 mm	$> 10^{4-6}$		$< 1 \text{ ms}$	10 ms	10	
<b>Intr. Signal</b>	10 $\mu\text{m}$	200 $\mu\text{m}$	$> 1000$		$< 1 \text{ ms}$	1 s	1000	10 ms
<b>VSD Imaging</b>	10 $\mu\text{m}$	30 $\mu\text{m}$	$< 100$		$< 1 \text{ ms}$	10 ms	$< 1$	
<b><math>\mu\text{Elec}</math></b>	50 $\mu\text{m}$	10 $\mu\text{m}$	1	10 $\mu\text{m}$	$< 1 \text{ ms}$	1 ms	1	
<b><math>\mu\text{Dialysis}</math></b>	$< 1 \text{ mm}$	100 $\mu\text{m}$	$> 10^4$		$> 1 \text{ min}$	100 ms	$> 10^6$	
<b>Intracell elec</b>	10 $\mu\text{m}$	10 $\mu\text{m}$	$< 1$		$< 1 \text{ ms}$	$< 1 \text{ ms}$	$< 1$	
<b>Ca imaging</b>	1 $\mu\text{m}$	1 $\mu\text{m}$	$< 1$		1 ms	$> 100 \text{ ms}$	$> 100$	

# Voltage Sensitive Dye (VSD)



Merocyanine  
Dye RH-890



# Voltage Sensitive Dye (VSD)

## MEASURES:


the weighted sum of membrane-potential changes in neuronal somata, dendritic and axonal arbors, and often glia

the dye signal is restricted to the site of the electrical activity

This signal mainly reflects the synaptic potentials in the dendrites

# Voltage Sensitive Dye (VSD)


In-vitro:



Modified from Cohen et al., 1972

# Voltage Sensitive Dye (VSD)


In-vivo:



Arieli, Lampl, Grinvald

# Voltage Sensitive Dye (VSD)


In-vivo:



# Voltage Sensitive Dye (VSD)

In-vivo:

Example: Surround Inhibition in the Rat Barrel Cortex



# Methods table

## Measuring neural activity

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
behavior	brain	brain	$> 10^{11}$	station	1 ms	10 ms	10	
2DG, c-fos	10 $\mu\text{m}$	10 $\mu\text{m}$	1		NA	30 min	$> 10^6$	
fMRI	$> 1$ mm	$> 200$ $\mu\text{m}$	$> 10^5$		1 s	$> 1$ s	1000	10 ms
EEG	10 mm	100 mm	$> 10^9$	station	$< 1$ ms	50 ms	50	1 ms
MEG	10 mm	100 mm	$> 10^9$		1 ms	50 ms	50	
ECoG	10 mm	1-10 mm	$> 10^{4-6}$		$< 1$ ms	10 ms	10	
Intr. Signal	10 $\mu\text{m}$	200 $\mu\text{m}$	$> 1000$		$< 1$ ms	1 s	1000	10 ms
VSD Imaging	10 $\mu\text{m}$	30 $\mu\text{m}$	$< 100$		$< 1$ ms	10 ms	$< 1$	
<b><math>\mu\text{Elec}</math></b>	<b>50 <math>\mu\text{m}</math></b>	<b>10 <math>\mu\text{m}</math></b>	<b>1</b>	<b>10 <math>\mu\text{m}</math></b>	<b><math>&lt; 1</math> ms</b>	<b>1 ms</b>	<b>1</b>	
$\mu\text{Dialysis}$	$< 1$ mm	100 $\mu\text{m}$	$> 10^4$		$> 1$ min	100 ms	$> 10^6$	
Intracell elec	10 $\mu\text{m}$	10 $\mu\text{m}$	$< 1$		$< 1$ ms	$< 1$ ms	$< 1$	
Ca imaging	1 $\mu\text{m}$	1 $\mu\text{m}$	$< 1$		1 ms	$> 100$ ms	$> 100$	

# Micro-electrode recordings



# Micro-electrode recordings

From the tip of the microelectrode one can record:


LFP – local field potential

MUA – multi-unit activity

SUA – single-unit activity (using spike sorting)




# Spikes, LFP and ECoG



# Micro-electrode recordings


## Multiple single-unit recordings using tetrodes



- Neurons are spike-sorted based on relative amplitudes on the 4 tetrode channels (amplitude differences are caused by physical proximity of neurons to different tetrode wires)
- Up to 25 cell can be well-separated per tetrode
- More typically: 5 – 10 cells per tetrode
- Can record > 100 neurons overall (in 10-20 tetrodes) in a freely behaving, freely moving animal

# Micro-electrode recordings

## Multiple single-unit recordings using tetrodes



- Here, 5 neurons were recorded simultaneously from one tetrode


# Methods table

## Measuring neural activity


	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>behavior</b>	brain	brain	$> 10^{11}$	station	1 ms	10 ms	10	
<b>2DG, c-fos</b>	10 $\mu\text{m}$	10 $\mu\text{m}$	1		NA	30 min	$> 10^6$	
<b>fMRI</b>	$> 1 \text{ mm}$	$> 200 \mu\text{m}$	$> 10^5$		1 s	$> 1 \text{ s}$	1000	10 ms
<b>EEG</b>	10 mm	100 mm	$> 10^9$	station	$< 1 \text{ ms}$	50 ms	50	1 ms
<b>MEG</b>	10 mm	100 mm	$> 10^9$		1 ms	50 ms	50	
<b>ECoG</b>	10 mm	1-10 mm	$> 10^{4-6}$		$< 1 \text{ ms}$	10 ms	10	
<b>Intr. Signal</b>	10 $\mu\text{m}$	200 $\mu\text{m}$	$> 1000$		$< 1 \text{ ms}$	1 s	1000	10 ms
<b>VSD Imaging</b>	10 $\mu\text{m}$	30 $\mu\text{m}$	$< 100$		$< 1 \text{ ms}$	10 ms	$< 1$	
<b><math>\mu\text{Elec}</math></b>	50 $\mu\text{m}$	10 $\mu\text{m}$	1	10 $\mu\text{m}$	$< 1 \text{ ms}$	1 ms	1	
<b><math>\mu\text{Dialysis}</math></b>	$< 1 \text{ mm}$	100 $\mu\text{m}$	$> 10^4$		$> 1 \text{ min}$	100 ms	$> 10^6$	
<b>Intracell elec</b>	10 $\mu\text{m}$	10 $\mu\text{m}$	$< 1$		$< 1 \text{ ms}$	$< 1 \text{ ms}$	$< 1$	
<b>Ca imaging</b>	1 $\mu\text{m}$	1 $\mu\text{m}$	$< 1$		1 ms	$> 100 \text{ ms}$	$> 100$	

# Micro-dialysis recordings

From the tip of the micro-dialysis probe one can record concentrations of chemicals



# Micro-dialysis recordings



# Methods table


## Measuring neural activity

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>behavior</b>	brain	brain	$> 10^{11}$	station	1 ms	10 ms	10	
<b>2DG, c-fos</b>	10 $\mu\text{m}$	10 $\mu\text{m}$	1		NA	30 min	$> 10^6$	
<b>fMRI</b>	$> 1 \text{ mm}$	$> 200 \mu\text{m}$	$> 10^5$		1 s	$> 1 \text{ s}$	1000	10 ms
<b>EEG</b>	10 mm	100 mm	$> 10^9$	station	$< 1 \text{ ms}$	50 ms	50	1 ms
<b>MEG</b>	10 mm	100 mm	$> 10^9$		1 ms	50 ms	50	
<b>ECoG</b>	10 mm	1-10 mm	$> 10^{4-6}$		$< 1 \text{ ms}$	10 ms	10	
<b>Intr. Signal</b>	10 $\mu\text{m}$	200 $\mu\text{m}$	$> 1000$		$< 1 \text{ ms}$	1 s	1000	10 ms
<b>VSD Imaging</b>	10 $\mu\text{m}$	30 $\mu\text{m}$	$< 100$		$< 1 \text{ ms}$	10 ms	$< 1$	
<b><math>\mu\text{Elec}</math></b>	50 $\mu\text{m}$	10 $\mu\text{m}$	1	10 $\mu\text{m}$	$< 1 \text{ ms}$	1 ms	1	
<b><math>\mu\text{Dialysis}</math></b>	$< 1 \text{ mm}$	100 $\mu\text{m}$	$> 10^4$		$> 1 \text{ min}$	100 ms	$> 10^6$	
<b>Intracell elec</b>	10 $\mu\text{m}$	10 $\mu\text{m}$	$< 1$		$< 1 \text{ ms}$	$< 1 \text{ ms}$	$< 1$	
<b>Ca imaging</b>	1 $\mu\text{m}$	1 $\mu\text{m}$	$< 1$		1 ms	$> 100 \text{ ms}$	$> 100$	

# Intra-cellular recordings

## In awake behaving mouse

- Identifying cell type
- Recording the inputs (reflecting individual inputs, network state, network activity)
- revealing mechanisms (ex-inh, ...)




C. Petersen and colleagues, 2010

# Methods table

## Measuring neural activity

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>behavior</b>	brain	brain	$> 10^{11}$	station	1 ms	10 ms	10	
<b>2DG, c-fos</b>	10 $\mu\text{m}$	10 $\mu\text{m}$	1		NA	30 min	$> 10^6$	
<b>fMRI</b>	$> 1 \text{ mm}$	$> 200 \mu\text{m}$	$> 10^5$		1 s	$> 1 \text{ s}$	1000	10 ms
<b>EEG</b>	10 mm	100 mm	$> 10^9$	station	$< 1 \text{ ms}$	50 ms	50	1 ms
<b>MEG</b>	10 mm	100 mm	$> 10^9$		1 ms	50 ms	50	
<b>ECoG</b>	10 mm	1-10 mm	$> 10^{4-6}$		$< 1 \text{ ms}$	10 ms	10	
<b>Intr. Signal</b>	10 $\mu\text{m}$	200 $\mu\text{m}$	$> 1000$		$< 1 \text{ ms}$	1 s	1000	10 ms
<b>VSD Imaging</b>	10 $\mu\text{m}$	30 $\mu\text{m}$	$< 100$		$< 1 \text{ ms}$	10 ms	$< 1$	
<b><math>\mu\text{Elec}</math></b>	50 $\mu\text{m}$	10 $\mu\text{m}$	1	10 $\mu\text{m}$	$< 1 \text{ ms}$	1 ms	1	
<b><math>\mu\text{Dialysis}</math></b>	$< 1 \text{ mm}$	100 $\mu\text{m}$	$> 10^4$		$> 1 \text{ min}$	100 ms	$> 10^6$	
<b>Intracell elec</b>	10 $\mu\text{m}$	10 $\mu\text{m}$	$< 1$		$< 1 \text{ ms}$	$< 1 \text{ ms}$	$< 1$	
<b>Ca imaging</b>	1 $\mu\text{m}$	1 $\mu\text{m}$	$< 1$		1 ms	$> 100 \text{ ms}$	$> 100$	


# 2P $\text{Ca}^{2+}$ imaging of RGCs in response to visual stimulation




- Left: Fluorescent image of RGCs expressing the  $\text{Ca}^{2+}$  indicator GCaMP6f.
- Right: Average  $\text{Ca}^{2+}$  transients in example RGCs in response to a 5 sec UV spot stimulation (represented by yellow bar).

# Ca imaging using 2-photon microscopy




Anesthetized mice



Ca




APs (juxtacell)



- Figure 5. Single-cell and population YC3.60 Ca<sup>2+</sup> signals in L2/3 of barrel cortex. (A)** Simultaneous two-photon Ca<sup>2+</sup> imaging in soma and dendrites of L2/3 neurons using vertical (xz-)imaging. Examples of spontaneous somatic (S, red) and apical dendritic (D, blue) YC3.60 Ca<sup>2+</sup> transients for the cells depicted in the left image. Right: Mean decay times in dendrites compared to somata for 23 measurements (gray lines; mean ± SEM). **(B)** Simultaneous juxtacellular voltage recording and two photon Ca<sup>2+</sup> imaging from a neuron showing rare events of sustained and high-frequency AP firing that are accompanied by large YC3.60 Ca<sup>2+</sup> transients with peak amplitudes of up to 30% ΔR/R. Top: Sustained AP firing leads to prolonged elevation of the fluorescence ratio. Bottom: A short burst of 11 APs is accompanied by a fast Ca<sup>2+</sup> transient, which returns to baseline following a stereotypical exponential decay. **(C)** Two-photon Ca<sup>2+</sup> imaging of a small population of neurons during sensory stimulation (seven times five air-puffs to contra-lateral whiskers at 5 Hz). Large Ca<sup>2+</sup> transients in cell 1 (red trace) correlated with the spiking activity observed in the simultaneous juxtacellular voltage recording. Concomitant Ca<sup>2+</sup> transients were also evoked in neighboring neuronal somata and in the nearby neuropil (NP). The response to the first stimulation episode (dashed box) is shown on expanded scale in the lower left, indicating that YC3.60 resolves the individual steps in the accumulated Ca<sup>2+</sup> response. **(D)** Event-triggered average Ca<sup>2+</sup> traces from somata and adjacent neuropil for spontaneous (*n* = 37 events of 1–3 APs) and evoked (*n* = 32 events of 1–5 APs) action potentials. Multi-whisker air puff evoked Ca<sup>2+</sup> transients in somata were significantly larger than those in the neuropil while spontaneous spikes were accompanied by somatic but no neuropil transients. Errors are shown as SEM.


# Ca imaging using 2-photon microscopy

Freely moving mice




- **Figure 7. Fiber-optic recording of brain area activity in freely moving mice using YC3.60.** (A,B) Two examples of fiber-optic recording of YC3.60 signals in awake, freely moving mice. Bulk  $\text{Ca}^{2+}$  signals indicating neuronal activity were recorded in somatosensory cortex through a single-core optical fiber as shown in Figure 6C. Fluorescence changes in the YFP-channel are shown during 25–30 s periods together with the position of the mouse in an open field box. Animal behavior (sitting still, moving, touches, or having contact to the wall) is indicated by background colors. The trajectory of the animals' movement is indicated with selected time stamps. (C) Six more examples of  $\text{Ca}^{2+}$  imaging from three mice, together with corresponding behavioral observations. Changes of the animal's behavioral state (e.g., start of movement) were frequently associated with marked discontinuities in the fluorescence trace, indicating complex underlying  $\text{Ca}^{2+}$  dynamics. (D) Control experiment showing that  $\text{Ca}^{2+}$  signals are blocked by local perfusion of the cortical region with  $\text{Cd}^{2+}$ . (E) Control experiment demonstrating that a flat fluorescence trace is observed in the absence of YC3.60 expression.

# A system for chronic imaging of neuronal ensemble activity in the hippocampus of freely behaving mice





# Imaging of $\text{Ca}^{2+}$ dynamics in CA1 neurons of freely behaving mice



# Ca imaging using 2-photon microscopy


Selecting only projecting neurons



- **Figure 1.** Retrograde labeling with a virus expressing a red fluorescent protein. **A**, Schematic showing retrograde labeling of neurons in SI by injection of the retrograde virus HSV1 into MI. **B**, *In vivo* image of RFP+ neurons (maximum-intensity side projection of an image stack of RFP+ neurons; 512 x 128 x 96; section spacing, 8  $\mu\text{m}$ ). **C**, Distribution of labeled neurons in SI barrel cortex after bead (left) or virus (right) injection into MI. The white lines indicate the pia and the border between the cortex and the white matter. **D**, Normalized distribution of labeled neurons after bead (black, 1293 neurons) and HSV (white, 808 neurons) injection into MI.

# Introduction

The scientific method



# Methods table

Stimulating / perturbing neural activity

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>sensory</b>	modality	modality	$> 10^9$	station	$< 1$ ms	$< 1$ ms	$< 1$	
<b>TMS</b>	100 mm	10 mm	$> 10^9$	station	$< 1$ ms	100 ms	100	
<b><math>\mu</math>Stim</b>	$< 10 \mu\text{m}$	$> 100 \mu\text{m}$	$> 50$		$< 1$ ms	10 ms	10	
<b><math>\mu</math>Pharmac</b>	$< 10 \mu\text{m}$	$> 100 \mu\text{m}$	$> 50$		1 ms	$> 10$ s	10	
<b>single cell</b>	$< 10 \mu\text{m}$	$< 10 \mu\text{m}$	1		$< 1$ ms	$< 1$ ms	$< 1$	
<b>sub-cell</b>	$> 100 \mu\text{m}$	$< 1 \mu\text{m}$	$> 50$		$< 1$ ms	$< 1$ ms	$< 1$	

# Methods table

Stimulating / perturbing neural activity


	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>sensory</b>	modality	modality	$> 10^9$	station	$< 1$ ms	$< 1$ ms	$< 1$	
<b>TMS</b>	100 mm	10 mm	$> 10^9$	station	$< 1$ ms	100 ms	100	
<b><math>\mu</math>Stim</b>	$< 10 \mu\text{m}$	$> 100 \mu\text{m}$	$> 50$		$< 1$ ms	10 ms	10	
<b><math>\mu</math>Pharmac</b>	$< 10 \mu\text{m}$	$> 100 \mu\text{m}$	$> 50$		1 ms	$> 10$ s	10	
<b>single cell</b>	$< 10 \mu\text{m}$	$< 10 \mu\text{m}$	1		$< 1$ ms	$< 1$ ms	$< 1$	
<b>sub-cell</b>	$> 100 \mu\text{m}$	$< 1 \mu\text{m}$	$> 50$		$< 1$ ms	$< 1$ ms	$< 1$	

# Methods table

Stimulating / perturbing neural activity

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>sensory</b>	modality	modality	$> 10^9$	station	$< 1$ ms	$< 1$ ms	$< 1$	
<b>TMS</b>	100 mm	10 mm	$> 10^9$	station	$< 1$ ms	100 ms	100	
<b><math>\mu</math>Stim</b>	$< 10 \mu\text{m}$	$> 100 \mu\text{m}$	$> 50$		$< 1$ ms	10 ms	10	
<b><math>\mu</math>Pharmac</b>	$< 10 \mu\text{m}$	$> 100 \mu\text{m}$	$> 50$		1 ms	$> 10$ s	10	
<b>single cell</b>	$< 10 \mu\text{m}$	$< 10 \mu\text{m}$	1		$< 1$ ms	$< 1$ ms	$< 1$	
<b>sub-cell</b>	$> 100 \mu\text{m}$	$< 1 \mu\text{m}$	$> 50$		$< 1$ ms	$< 1$ ms	$< 1$	


# Micro-stimulation



E Seidemann et al. Science 2002;295:862-865

Figure 1 Spatiotemporal dynamics of microstimulation-evoked activity.

# Micro-stimulation



E Seidemann et al. Science 2002;295:862-865

Figure 2 Time course of the response to microstimulation.

# Methods table

Stimulating / perturbing neural activity

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>sensory</b>	modality	modality	$> 10^9$	station	$< 1$ ms	$< 1$ ms	$< 1$	
<b>TMS</b>	100 mm	10 mm	$> 10^9$	station	$< 1$ ms	100 ms	100	
<b><math>\mu</math>Stim</b>	$< 10 \mu\text{m}$	$> 100 \mu\text{m}$	$> 50$		$< 1$ ms	10 ms	10	
<b><math>\mu</math>Pharmac</b>	$< 10 \mu\text{m}$	$> 100 \mu\text{m}$	$> 50$		1 ms	$> 10$ s	10	
<b>single cell</b>	$< 10 \mu\text{m}$	$< 10 \mu\text{m}$	1		$< 1$ ms	$< 1$ ms	$< 1$	
<b>sub-cell</b>	$> 100 \mu\text{m}$	$< 1 \mu\text{m}$	$> 50$		$< 1$ ms	$< 1$ ms	$< 1$	





# Micro-pharmacology

combined electrode:

Extracellular recording &  
drug application

Inject

- charged chemicals using **iontophoresis**
- uncharged chemicals using **pressure**



Haidarliu, et al 1995

# Pharmacogenetics

- PSAM: a chimeric ligand-gated chloride channel. The channels are reversibly opened by the synthetic ligand PSEM.
- Here, PSAM was genetically expressed in one type of retinal neurons: starburst amacrine cells. These neurons are thought to mediate direction selective responses in the retina.

- Left: Two-photon fluorescence image of the ganglion cell layer of a retina expressing PSAM. Circles are cells identified as direction selective ganglion cells.
- Right: Example tuning curve of a direction selective ganglion cell before (Control), during (PSEM) and after (Wash) the addition of PSEM. Direction selective responses are abolished when starburst cells are inhibited.

# Methods table



Stimulating / perturbing neural activity

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>sensory</b>	modality	modality	$> 10^9$	station	$< 1$ ms	$< 1$ ms	$< 1$	
<b>TMS</b>	100 mm	10 mm	$> 10^9$	station	$< 1$ ms	100 ms	100	
<b><math>\mu</math>Stim</b>	$< 10 \mu\text{m}$	$> 100 \mu\text{m}$	$> 50$		$< 1$ ms	10 ms	10	
<b><math>\mu</math>Pharmac</b>	$< 10 \mu\text{m}$	$> 100 \mu\text{m}$	$> 50$		1 ms	$> 10$ s	10	
<b>single cell</b>	$< 10 \mu\text{m}$	$< 10 \mu\text{m}$	1		$< 1$ ms	$< 1$ ms	$< 1$	
<b>sub-cell</b>	$> 100 \mu\text{m}$	$< 1 \mu\text{m}$	$> 50$		$< 1$ ms	$< 1$ ms	$< 1$	

# Nano-stimulation



Examples of whisker protraction and retraction caused by single and multiple spikes

Anesthetized rats




Herfst, L. J. et al. *J Neurophysiol* 99: 2821-2832 2008;  
doi:10.1152/jn.01014.2007

# Nano-stimulation



Behaving rats



Houweling & Michael Brecht, Nature 2008

# Methods table


Stimulating / perturbing neural activity

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>sensory</b>	modality	modality	$> 10^9$	station	$< 1$ ms	$< 1$ ms	$< 1$	
<b>TMS</b>	100 mm	10 mm	$> 10^9$	station	$< 1$ ms	100 ms	100	
<b><math>\mu</math>Stim</b>	$< 10 \mu\text{m}$	$> 100 \mu\text{m}$	$> 50$		$< 1$ ms	10 ms	10	
<b><math>\mu</math>Pharmac</b>	$< 10 \mu\text{m}$	$> 100 \mu\text{m}$	$> 50$		1 ms	$> 10$ s	10	
<b>single cell</b>	$< 10 \mu\text{m}$	$< 10 \mu\text{m}$	1		$< 1$ ms	$< 1$ ms	$< 1$	
<b>sub-cell</b>	$> 100 \mu\text{m}$	$< 1 \mu\text{m}$	$> 50$		$< 1$ ms	$< 1$ ms	$< 1$	


# Optogenetic-stimulation

## Channelrhodopsins (ChR1,2)

- Chrs function as light-gated ion channels.
- They serve as sensory photoreceptors in unicellular green algae, controlling phototaxis.
- Expressed in cells of other organisms, they enable the use of light to control electrical excitability
- All known Chrs are nonspecific cation channels, conducting  $H^+$ ,  $Na^+$ ,  $K^+$ , and  $Ca^{2+}$  ions.






Cultured neuron expressing ChR2-eYFP stimulated with pulses of blue light



# Optogenetic-stimulation

- Variety of excitatory channels: ON short duration, ON long duration, ON/OFF, ON subliminal, ...
- Inhibitory channels
- Linking to identified promoters to be functional in identified neurons
- Localized infection using viruses
- Combined with dye reporters








# Methods table


## Measuring structure

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
cell density								
receptor density								
transmitter density								
tract tracing								
single-cell								
single-spine	< 1 $\mu\text{m}$	< 1 $\mu\text{m}$	< 1		< 1 s	hours		



# Single-spine monitoring



# Single-spine monitoring



# Single-spine monitoring



# Methods table

## Manipulating structure

	Spatial resolution				Temporal resolution			
	<u>device</u>	<u>signal</u>	<u>neurons</u>	<u>heuristic</u>	<u>device</u>	<u>signal</u>	<u>spikes</u>	<u>heuristic</u>
<b>Neuropsychology</b>		> 10 mm	> 10 <sup>7</sup>			months		
<b>lesions</b>	> 100 μm	> 100 μm	> 50		> 1 s	> 1 min		