

previously shown to be required for EutV binding of dual-hairpin substrates (5). We tested one mutant, in which these critical residues of the P3 loop were mutated (M1) and a second, in which the same changes were made to the P4 loop (M2). Similar to deletion of P3/P4, these mutant constructs were both induced by EA alone in the *eutX* strain, in contrast to wild-type *eutG-lacZ* (Fig. 3D). Therefore, these data confirm that P3/P4 mutations that prevent EutV binding affect regulation. We also examined the M3 mutation predicted to abrogate AdoCbl binding and cause constitutive inclusion of P3/P4 in EutX (Fig. 2, C and D). As expected, this resulted in uninducible expression (Fig. 3C).

In conclusion, we present evidence of a riboswitch acting in trans, the second described (13). In the prior example, a classical sRNA that acts by base-pairing to a target mRNA can be prematurely terminated by a riboswitch. However, EutX acts in a signal transduction pathway that is subject to riboswitch-mediated control within the confines of an sRNA. A few other sRNAs, like EutX, have been shown to affect gene expression by sequestering RNA-binding proteins. The best-studied examples are CsrB and CsrC, which titrate CsrA, a translational inhibitor protein, away from its mRNA targets [reviewed by (14)]. EutX demonstrates a mechanism for how protein sequestration can be placed under signal-responsive regulatory control by a riboswitch.

REFERENCES AND NOTES

1. A. Roth, R. R. Breaker, *Annu. Rev. Biochem.* **78**, 305–334 (2009).
2. A. Serganov, E. Nudler, *Cell* **152**, 17–24 (2013).
3. L. S. Waters, G. Storz, *Cell* **136**, 615–628 (2009).
4. K. A. Fox et al., *Proc. Natl. Acad. Sci. U.S.A.* **106**, 4435–4440 (2009).
5. A. Ramesh et al., *PLOS Genet.* **8**, e1002666 (2012).
6. F. A. Scarlett, J. M. Turner, *J. Gen. Microbiol.* **95**, 173–176 (1976).
7. D. A. Garsin, *Nat. Rev. Microbiol.* **8**, 290–295 (2010).
8. K. A. Baker, M. Perego, *J. Bacteriol.* **193**, 2575–2586 (2011).
9. M. F. Del Papa, M. Perego, *J. Bacteriol.* **190**, 7147–7156 (2008).
10. J. E. Johnson Jr., F. E. Reyes, J. T. Polaski, R. T. Batey, *Nature* **492**, 133–137 (2012).
11. A. Pesellis, A. Serganov, *Nat. Struct. Mol. Biol.* **19**, 1182–1184 (2012).
12. K. G. Roelofs, J. Wang, H. O. Sintim, V. T. Lee, *Proc. Natl. Acad. Sci. U.S.A.* **108**, 15528–15533 (2011).
13. E. Loh et al., *Cell* **139**, 770–779 (2009).
14. P. Babinzke, T. Romeo, *Curr. Opin. Microbiol.* **10**, 156–163 (2007).

ACKNOWLEDGMENTS

The data reported are presented in the main paper and the supplementary materials. This work was supported by the NIH grants RO1A076406 (D.A.G.), R56A110432 (D.A.G. and W.C.W.) and RO1GM099790 (A.V.H.), a NSF grant MCB-1051440 (W.C.W.) and a Welch Foundation grant AU-1773 (A.V.H.). For transmission electron microscopy imaging, we thank S. Kolodziej and P. Navarro; J. R. Mellin and P. Cossart are acknowledged for valuable discussions.

SUPPLEMENTARY MATERIALS

www.sciencemag.org/content/345/6199/937/suppl/DC1
Materials and Methods
Figs. S1 to S7
Tables S1 and S2
References (15–17)

21 April 2014; accepted 26 June 2014
10.1126/science.1255091

RIBOSWITCHES

Sequestration of a two-component response regulator by a riboswitch-regulated noncoding RNA

J. R. Mellin,^{1,2,3} Mikael Koutero,^{1,2,3} Daniel Dar,⁴ Marie-Anne Nahori,^{1,2,3} Rotem Sorek,⁴ Pascale Cossart^{1,2,3*}

Riboswitches are ligand-binding elements contained within the 5' untranslated regions of bacterial transcripts, which generally regulate expression of downstream open reading frames. Here, we show that in *Listeria monocytogenes*, a riboswitch that binds vitamin B₁₂ controls expression of a noncoding regulatory RNA, Rli55. Rli55, in turn, controls expression of the *eut* genes, whose products enable ethanolamine utilization and require B₁₂ as a cofactor. Defects in ethanolamine utilization, or in its regulation by Rli55, significantly attenuate *Listeria* virulence in mice. Rli55 functions by sequestering the two-component response regulator EutV by means of a EutV-binding site contained within the RNA. Thus, Rli55 is a riboswitch-regulated member of the small group of regulatory RNAs that function by sequestering a protein and reveals a distinctive mechanism of signal integration in bacterial gene regulation.

Ethanolamine is an abundant molecule in the vertebrate intestine (1, 2), and genes of the ethanolamine utilization pathway (*eut*) are widely conserved in pathogenic bacteria (3–5). This includes the Gram-positive intracellular human pathogen *Listeria monocytogenes*, in which *eut* expression has been shown to be up-regulated in the intestine during infection of mice (6), which suggests that ethanolamine is important for *Listeria* pathogenesis. In *Enterococcus faecalis*, *eut* expression is activated in response to ethanolamine by a two-component response regulator, EutVW (7, 8). In *Salmonella enterica*, ethanolamine utilization requires vitamin B₁₂ as a cofactor (9), and we noted the presence of a B₁₂-binding riboswitch located upstream of the first gene in the *eut* locus of *L. monocytogenes* (Fig. 1A) (10), which suggested that *eut* expression might also be regulated in response to B₁₂ availability.

To investigate a role for B₁₂, we examined expression of the *eut* locus in response to B₁₂ and ethanolamine by RNA sequencing (RNA-seq) (Fig. 1B) and quantitative reverse transcription polymerase chain reaction (qRT-PCR) (fig. S1). We observed expression of the *eutVW* genes under all conditions, albeit at low levels, which suggested that the cell maintains a pool of EutVW to sense and respond to ethanolamine. In contrast, higher-level expression of *eutVW* and expression of other *eut* genes require both B₁₂ and ethanolamine (Fig. 1B). These data indicated B₁₂ is required to activate *eut* expression and suggested that the B₁₂ riboswitch does not prevent

transcription of the *eut* locus in the presence of B₁₂, as might be expected for a classical riboswitch (11).

To clarify whether the riboswitch has a role in the B₁₂-dependent regulation of *eut* expression, we examined transcription of the riboswitch locus (Fig. 1C) in response to B₁₂ and ethanolamine. We were unable to detect any long transcript, which might extend into the downstream *eut* locus. However, we did detect a ~450-nucleotide (nt) transcript, Rli55 (10, 12), in the absence of B₁₂ (Fig. 1D), and a smaller (~200-nt) transcript that accumulated in the presence of B₁₂ (Fig. 1D). This result suggested that the riboswitch mediates transcription termination of the ~450-nt Rli55 transcript in response to B₁₂, and any remaining long transcript is rapidly degraded (fig. S2A). A strain with a deletion in the B₁₂ riboswitch (Δ ribo) constitutively expressed the long Rli55 transcript under all conditions (Fig. 1E and fig. S2B), which confirmed B₁₂-dependent regulation by the riboswitch. Conversely, neither the long or short Rli55 transcripts were detected in a strain (Δ rli55) in which the entire *rli55* locus was deleted (Fig. 1E and fig. S2B). Thus, the B₁₂ riboswitch determines whether Rli55 is expressed as a long or short transcript in response to B₁₂.

We reasoned that Rli55 might act as a regulatory RNA controlling expression of the *eut* locus. To test this hypothesis, we examined expression of *eut* genes by qRT-PCR in the wild-type, Δ ribo, and Δ rli55 strains. In the wild-type strain, *eut* expression was elevated only when both ethanolamine and B₁₂ were present in the media (Fig. 1F and fig. S3, A and B), whereas in the Δ ribo strain, which constitutively expresses the long Rli55, *eut* genes were never expressed under any conditions. In contrast, in the Δ rli55 strain, expression of the *eut* genes was high in the presence of ethanolamine alone in addition

¹Unité des Interactions Bactéries-Cellules, Institut Pasteur, F-75015 Paris, France. ²INSERM, U604, Paris, F-75015 France.

³INRA, USC2020, F-75015 Paris, France. ⁴Department of Molecular Genetics, Weizmann Institute of Science, Rehovot, Israel.

*Corresponding author. E-mail: pcossart@pasteur.fr





Fig. 2. Ethanolamine utilization and pathogenesis. BALB/c mice were injected intravenously with ~4500 colony-forming units (CFU) of the indicated strain. Mice were killed at (A) and (D) 24 hours, (B) and (E) 48 hours, or (C) and (F) 72 hours, and spleens and livers were removed to assess bacterial load per organ. Results represent two independent experiments with three or four mice per group in each experiment. * $P < 0.05$, ns, not significant.



to ethanolamine and B₁₂ together. Together, these data support a model in which Rli55 prevents the expression of the *eut* locus in the absence of B₁₂, ensuring that the *eut* genes are expressed only in the presence of both ethanolamine (substrate) and B₁₂ (cofactor) (fig. S3C).

We tested whether defects in ethanolamine utilization, or its regulation, impacted *L. monocytogenes* virulence by examining the wild-type, Δ ribo, and Δ rli55 strains in a mouse intravenous infection model. We also tested a mutant lacking the *eutB* gene encoding an ethanolamine lyase subunit, which is unable to catabolize ethanolamine. The Δ ribo and Δ eutB strains both had significantly reduced bacterial loads at 24 hours after infection compared with the wild-type strain, and these differences increased at 48 hours and 72 hours postinfection (~10- to 50-fold) (Fig. 2, A to F). In contrast, the Δ rli55 strain, in which *eut* expression is not inhibited, was present in amounts comparable to those in the wild-type strain in the spleen and liver. Thus,

defects in ethanolamine utilization or activation of *eut* expression significantly attenuate *L. monocytogenes* virulence.

In *E. faecalis*, ethanolamine is sensed by the sensor-kinase EutW, which subsequently phosphorylates the response regulator EutV (7). Phosphorylated EutV in turn binds ANTAR elements (AmiR and NasR transcriptional antiterminator regulator) in the 5' untranslated regions of actively transcribed *eut* mRNAs, which prevents the formation of a transcription terminator and consequently activates *eut* expression (13). In *L. monocytogenes*, ANTAR sites were identified upstream of the *eutA* and *eutV* genes, and a third site was identified in the *rli55* locus upstream of the *eutG* gene and downstream of the B₁₂ riboswitch (13). We also identified a second ANTAR site in the *rli55* locus (Fig. 3A), which suggested that one or both of these ANTAR elements could be transcribed as part of the 3' end of Rli55 RNA. Indeed, our RNA-seq data showed that, in the absence of B₁₂, high levels of Rli55

are transcribed as a long transcript encompassing the first ANTAR element (Fig. 3A). In contrast, *rli55* transcription terminates abruptly after the riboswitch in the presence of B₁₂, which indicates that the riboswitch determines whether Rli55 is transcribed with or without an ANTAR element. This notion was supported by the detection of putative Rli55 orthologs in *E. faecalis* and *Streptococcus sanguinis* (fig. S4). In the latter, the riboswitch, in conjunction with a single ANTAR element, has undergone an inversion relative to the adjacent *eutG* gene, which suggests that the riboswitch and the first ANTAR element are functionally linked (fig. S4B) and that this is also the case in *L. monocytogenes*.

To test if the ANTAR elements are involved in Rli55-mediated regulation, we complemented the Δ rli55 strain with chromosomally integrated *rli55* alleles carrying mutations in the ANTAR elements and examined which ones restored Rli55-mediated regulation of *eut* expression (Fig. 3, B and C, and fig. S5). A strain with an empty




Fig. 3. Rli55 sequesters the two-component response regulator EutV.

(A) RNA-seq coverage in reads per million (RPM) of the *rli55* locus from bacteria grown as indicated. (B and C) qRT-PCR of *eutA* and *eutV* genes. Vitamin B₁₂ (B₁₂), ethanolamine (Ea). Values represent means \pm SEM, $n = 3$. All differences were significantly different (* $P < 0.05$) from the Δ rli55::Empty strain in the Ea condition, except where indicated (ns). All differences were not significantly different from the Δ rli55::Empty strain in the B₁₂ + Ea condition, ex-

cept where indicated. (D) Secondary structure of the ANTAR elements encoded in the boxed region of Fig. 3A. Mutations are shown in blue. (E and F) RNA-seq coverage in RPM of the *rli55* and *eutV* loci with RNA isolated by coimmunoprecipitation of cell lysates from either EutV^{FLAG} or EutV^{NOFLAG} cultures grown in the presence of ethanolamine (Ea) or ethanolamine and B₁₂ together. (G) Proposed model of Rli55-mediated regulation of *eut* expression in the presence of ethanolamine (Ea) alone or ethanolamine + B₁₂ (Ea + B₁₂).

construct ($\Delta rli55::\text{Empty}$) could not prevent expression of the *eut* genes in ethanolamine alone (as in the parental $\Delta rli55$ strain), whereas a strain with a wild-type copy of *rli55* ($\Delta rli55::rli55$) fully restored Rli55-mediated inhibition. However, a strain with a deletion in the riboswitch ($\Delta rli55::\Delta ribo$) inhibited *eut* expression in all conditions, as the riboswitch can no longer terminate *rli55* transcription in response to B_{12} . In strain $\Delta rli55::rli55\Delta M1$, wherein four uridine residues in the first ANTAR site were mutated to adenines (Fig. 3D, $\Delta M1$), inhibition of *eut* expression by Rli55 was abolished in the presence of ethanolamine alone. In contrast, in strain $\Delta rli55::rli55\Delta M1/M2$, where compensatory mutations were made to the opposite side of the ANTAR stem-loop (Fig. 3D, $\Delta M2$), wild-type regulation of *eut* expression was restored. Mutation of the six nucleotides in the stem-loop of the second ANTAR element (Fig. 3D, $\Delta M3$, $\Delta rli55::rli55\Delta M3$) had no significant effect on Rli55-mediated regulation. Thus, the first ANTAR element is necessary and sufficient for Rli55-mediated regulation.

The long form of Rli55 containing an ANTAR element might bind and sequester EutV and so prevent it from activating expression of the *eut* genes in the presence of ethanolamine but absence of B_{12} . When sufficient levels of B_{12} accumulate, B_{12} would bind the riboswitch, producing truncated Rli55 transcripts, which would lack an ANTAR element and be unable to sequester EutV. To examine this hypothesis, we constructed a strain with an additional copy of the *eutV* gene carrying a 2XFLAG-tag ($\text{EutV}^{\text{FLAG}}$) and first showed that expression of $\text{EutV}^{\text{FLAG}}$ protein is regulated identically to the native *eutV* gene in response to ethanolamine and B_{12} (fig. S6A). We also constructed a strain with an additional *eutV* gene lacking a FLAG tag ($\text{EutV}^{\text{NOFLAG}}$). Anti-FLAG immunoprecipitations of cell lysates from these two strains (fig. S6, B and C), followed by RNA-seq analysis (Fig. 3, E and F), showed that Rli55 is enriched by co-immunoprecipitation with $\text{EutV}^{\text{FLAG}}$ primarily when bacteria are grown in the presence of ethanolamine alone, although we saw no enrichment in a parallel immunoprecipitation with the $\text{EutV}^{\text{NOFLAG}}$ strain (Fig. 3E). In contrast, the ANTAR element upstream of the *eutV* gene (Fig. 3F) is enriched by coimmunoprecipitation of lysates from $\text{EutV}^{\text{FLAG}}$ bacteria, but not $\text{EutV}^{\text{NOFLAG}}$ bacteria, grown in the presence of ethanolamine and B_{12} together but not from lysates of bacteria grown in ethanolamine alone. To a lesser extent, the ANTAR-containing region upstream of *eutA* and the entire *eutA-Q* locus are enriched under the latter condition (fig. S7). These data support a model in which the majority of EutV is bound and sequestered by Rli55 in the presence of ethanolamine alone. Conversely, in the presence of ethanolamine and B_{12} , the riboswitch produces short truncated Rli55 transcripts, which cannot bind EutV, and so allows EutV to bind *eut* mRNAs and to activate *eut* expression (Fig. 3G).

This riboregulatory mechanism coordinates expression of the ethanolamine utilization (*eut*)

locus with the availability of B_{12} , the essential cofactor for ethanolamine catabolism. Previously, ethanolamine utilization has been shown to be important after oral infection by *Salmonella enterica* serovar Typhimurium and enterohemorrhagic *Escherichia coli* (3, 4, 14); however, the contribution of ethanolamine utilization to *L. monocytogenes* pathogenesis in an intravenous mouse infection model suggests that ethanolamine utilization is important outside of the intestine and possibly in the intracellular environment. This study also extends the role of riboswitches in the regulation of noncoding RNAs (15, 16). Finally, our data show that Rli55 represents a new member of the small family of regulatory RNAs that function by sequestering a protein, which also includes the 6S and CsrB/C RNAs (17), and highlights a distinctive means of signal integration in bacterial gene regulation.

REFERENCES AND NOTES

- D. A. Garsin, *Nat. Rev. Microbiol.* **8**, 290–295 (2010).
- K. Koichi, F. Michiya, N. Makoto, *Biochim. Biophys. Acta* **369**, 222–233 (1974).
- P. Thiennimitr et al., *Proc. Natl. Acad. Sci. U.S.A.* **108**, 17480–17485 (2011).
- Y. Bertin et al., *Environ. Microbiol.* **13**, 365–377 (2011).
- O. Tsoy, D. Ravcheev, A. Mushegian, *J. Bacteriol.* **191**, 7157–7164 (2009).
- C. Archambaud et al., *Proc. Natl. Acad. Sci. U.S.A.* **109**, 16684–16689 (2012).
- M. F. Del Papa, M. Perego, *J. Bacteriol.* **190**, 7147–7156 (2008).
- K. A. Fox et al., *Proc. Natl. Acad. Sci. U.S.A.* **106**, 4435–4440 (2009).
- J. R. Roth, J. G. Lawrence, T. A. Bobik, *Annu. Rev. Microbiol.* **50**, 137–181 (1996).
- A. Toledo-Arana et al., *Nature* **459**, 950–956 (2009).
- W. C. Winkler, R. R. Breaker, *Annu. Rev. Microbiol.* **59**, 487–517 (2005).
- O. Wurtzel et al., *Mol. Syst. Biol.* **8**, 583 (2012).
- A. Ramesh et al., *PLoS Genet.* **8**, e1002666 (2012).
- S. E. Winter, A. J. Bäumer, *Gut Microbes* **2**, 58–60 (2011).
- J. R. Mellin et al., *Proc. Natl. Acad. Sci. U.S.A.* **110**, 13132–13137 (2013).
- G. André et al., *Nucleic Acids Res.* **36**, 5955–5969 (2008).
- S. Gottesman, G. Storz, *Cold Spring Harb. Perspect. Biol.* **3**, a003798 (2011).

ACKNOWLEDGMENTS

We are grateful to D. Garsin and W. Winkler for their exchange of ideas and discussion of unpublished data. We also thank N. King and J. Pederson for helpful and insightful discussions and Y. Chao and J. Vogel for assistance with the CLIP-seq protocol. This work was supported by grants to P.C., European Research Council advanced grant (233348), ANR (BACNET 09-BLAN-0024-02), ANR Investissement d'Avenir Programme (10-LABX-62), Fondation Le Roch, and Fondation Jeantet. P.C. is an International Senior Research Scholar of the Howard Hughes Medical Institute.

SUPPLEMENTARY MATERIALS

www.sciencemag.org/content/345/6199/940/suppl/DC1
Materials and Methods
Figs. S1 to S7
Tables S1 and S2
References

21 April 2014; accepted 24 June 2014
10.1126/science.1255083

IMMUNOGENETICS

Chromatin state dynamics during blood formation

David Lara-Astiaso,^{1,*} Assaf Weiner,^{2,3,*} Erika Lorenzo-Vivas,¹ Irina Zaretsky,¹ Diego Adhemar Jaitin,¹ Eyal David,¹ Hadas Keren-Shaul,¹ Alexander Mildner,¹ Deborah Winter,¹ Steffen Jung,¹ Nir Friedman,^{2,3,†} Ido Amit^{1,†}

Chromatin modifications are crucial for development, yet little is known about their dynamics during differentiation. Hematopoiesis provides a well-defined model to study chromatin state dynamics; however, technical limitations impede profiling of homogeneous differentiation intermediates. We developed a high-sensitivity indexing-first chromatin immunoprecipitation approach to profile the dynamics of four chromatin modifications across 16 stages of hematopoietic differentiation. We identify 48,415 enhancer regions and characterize their dynamics. We find that lineage commitment involves de novo establishment of 17,035 lineage-specific enhancers. These enhancer repertoire expansions foreshadow transcriptional programs in differentiated cells. Combining our enhancer catalog with gene expression profiles, we elucidate the transcription factor network controlling chromatin dynamics and lineage specification in hematopoiesis. Together, our results provide a comprehensive model of chromatin dynamics during development.

Chromatin plays a major regulatory role in cell-type-specific functions and response (1, 2). The current dogma of cellular differentiation suggests that there is a progressive closing of the regulatory potential of the genome. According to this model, differentia-

tion is a gradual transition from an open chromatin state in multipotent stem cells to a compacted chromatin state in differentiated cells. However, genome-wide histone modification profiling of embryonic stem cells and terminally differentiated cells is not fully compatible with this model

Sequestration of a two-component response regulator by a riboswitch-regulated noncoding RNA

J. R. Mellin, Mikael Koutero, Daniel Dar, Marie-Anne Nahori, Rotem Sorek and Pascale Cossart

Science **345** (6199), 940-943.
DOI: 10.1126/science.1255083

A dual-action RNA switch for expression

Riboswitches are short segments of RNA that bind small molecules and switch between two different conformations, thereby regulating gene expression (see the Perspective by Chen and Gottesman). DebRoy *et al.* and Mellin *et al.* find a new class of riboswitches—in two different species of bacteria—that are both part of and regulate the production of a noncoding RNA. Each riboswitch ensures that a particular metabolic pathway is only activated in the presence of an essential small-molecule cofactor. In the absence of the cofactor, the full-length non-coding RNA is made and binds a regulator protein, preventing the regulator protein from inappropriately activating the metabolic pathway.

Science, this issue p. 937 and p. 940; see also p. 876

ARTICLE TOOLS	http://science.sciencemag.org/content/345/6199/940
SUPPLEMENTARY MATERIALS	http://science.sciencemag.org/content/suppl/2014/08/20/345.6199.940.DC1
RELATED CONTENT	http://science.sciencemag.org/content/sci/345/6199/876.full http://science.sciencemag.org/content/sci/345/6199/937.full
REFERENCES	This article cites 17 articles, 7 of which you can access for free http://science.sciencemag.org/content/345/6199/940#BIBL
PERMISSIONS	http://www.sciencemag.org/help/reprints-and-permissions

Use of this article is subject to the [Terms of Service](#)

Science (print ISSN 0036-8075; online ISSN 1095-9203) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science* is a registered trademark of AAAS.

Copyright © 2014, American Association for the Advancement of Science